


PCT

WORLD INTELLECTUAL PROPRIETY  
International B



INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 9607739A2

(51) International Patent Classification 6: C12N 15/12, C07K 14/715, 16/28, A61K 38/17, C12Q 1/68	A2	(11) International Publication Number: WO 96/07739 (43) International Publication Date: 14 March 1996 (14.03.96)
---	----	---

(21) International Application Number: PCT/US95/12037

(22) International Filing Date: 11 September 1995 (11.09.95)

(30) Priority Data:  
08/303,957 9 September 1994 (09.09.94) US

(71) Applicant: NEUROCRINE BIOSCIENCES, INCORPORATED [US/US]; 3050 Science Park Road, San Diego, CA 92121 (US).

(72) Inventors: LOVENBERG, Timothy, W.; 6529 Vispera Place, Carlsbad, CA 92009 (US). OLTERSDORF, Tilman; 427 Bristol Avenue, Cardiff, CA 92007 (US). LIAW, Chen, W.; 7668 Salix Place, San Diego, CA 92129 (US). CLEVELINGER, William; 1510 South Melrose Drive #7, Vista, CA 92083 (US). DeSOUZA, Errol, B.; 4507 South Lane, Del Mar, CA 92014 (US).

(74) Agents: McMASTERS, David, D. et al.; Seed and Berry, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).

(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).


Published

Without international search report and to be republished upon receipt of that report.

(54) Title: INTERLEUKIN-1 TYPE 3 RECEPTORS

(57) Abstract

The present invention provides isolated nucleic acid molecules encoding soluble and membrane bound forms of Interleukin-1 Type 3 receptors, as well as recombinant expression vectors and host cells suitable for expressing such receptors.



RAT IL-1 TYPE 3 RECEPTOR

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## Description

### INTERLEUKIN-1 TYPE 3 RECEPTORS

#### 5 Technical Field

The present invention relates generally to cell surface receptors, and more specifically, to Interleukin-1 Type 3 receptors.

#### Background Of The Invention

10 Interleukin-1 ("IL-1") is a cytokine which is known to be a key mediator of immunological and pathological responses to stress, infection and antigenic challenge (Oppenheim et al., *Immunol. Today* 7:45-46, 1986; Dinarello, *FASEB J.* 2:108-115, 1988; and Mizel, *FASEB J.* 3:2379-2388, 1989). In addition, IL-1 is known to have a variety of effects on the brain and central nervous system. For example, IL-1 has been  
15 postulated to be involved in the induction of fever (Kluger, *Physiol. Rev.* 71:93-127, 1991), increased duration of slow wave sleep (Opp et al., *Am. J. Physiol.* 260:R52-R58, 1991), decreased appetite (McCarthy et al., *Am. J. Clin. Nutr.* 42:1179-1182, 1985), activation of the hypothalamic-pituitary-adrenal ("HPA") axis (Woloski et al., *Science* 230:1035-1037, 1985), and inhibition of the hypothalamic-pituitary-gonadal axis (River  
20 and Vale, *Endocrinology* 124:2105-2109, 1989).

In light of the above-noted effects of IL-1 (as well as many others), substantial effort has been undertaken in order to identify receptors for IL-1. Briefly, at least two types of receptors are known to be expressed on the surface of certain immune cells in both human and murine derived lines. Type I receptors bind both IL-1 $\alpha$  and  
25 IL-1 $\beta$ , and can be found on T cells, fibroblasts, keratinocytes, endothelial cells, synovial lining cells, chondrocytes and hepatocytes (U.S. Patent Nos. 4,968,607, 5,081,228, and 5,180,812; Chizzonite et al., *PNAS* 86:8029-8033, 1989; Dinarello et al., *Blood* 77:1627-1652, 1991). Type II receptors can be found on various B cell lines, including the Raji human B-cell lymphoma line (Bomsztyk et al., *PNAS* 86:8034-8038, 1989;  
30 Horuk et al., *J. Biol. Chem.* 262:16275-16278, 1987; Horuk and McCubrey, *Biochem. J.* 260:657-663, 1989).

The present invention provides new, previously unidentified Interleukin receptors, designated Interleukin-1 Type 3 receptors ("IL-1-3R"). In addition, the present invention provides compositions and methods which utilize such receptors, as  
35 well as other, related advantages.

Summary of the Invention

Briefly stated, the present invention provides compositions and methods which comprise Interleukin-1 Type 3 receptors. Within one aspect of the present invention isolated nucleic acid molecules are provided which encode Interleukin-1 Type 3 receptors. Within one embodiment, the isolated nucleic acid molecules comprise the sequence of nucleotides in Sequence I.D. No. 1, from nucleotide number 129 to nucleotide number 1814. Within another embodiment, the isolated nucleic acid molecules encode a protein having the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 562. Within other embodiments, isolated nucleic acid molecules are provided in Sequence I.D. No. 3, from nucleotide number 89 to nucleotide number 1771. Within another embodiment, the nucleic acid molecules encode a protein having the amino acid sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 561. Nucleic acid molecules which encode IL-1 Type 3 receptors of the present invention may be isolated from virtually any warm-blooded animal, including for example, humans, macaques, horses, cattle, sheep, pigs, dogs, cats, rats and mice.

Within related aspects of the present invention, isolated nucleic acid molecules are provided which encode soluble Interleukin-1 Type 3 receptors. Within one embodiment, the isolated nucleic acid molecules comprise the sequence of nucleotides in Sequence I.D. No. 1, from nucleotide number 129 to nucleotide number 1136. Within other embodiments, the isolated nucleic acid molecules encode a protein having the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 336. Within another embodiment, the nucleic acid molecules comprise the sequence of nucleotides in Sequence I.D. No. 3, from nucleotide number 89 to nucleotide number 1102. Within yet another embodiment, the nucleic acid molecules encode a protein having the amino acid sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 338. As above, nucleic acid molecules which encode soluble IL-1 Type 3 receptors of the present invention may be isolated from virtually any warm-blooded animal, including for example, humans, macaques, horses, cattle, sheep, pigs, dogs, cats, rats and mice.

Within other aspects of the present invention, expression vectors are provided which are capable of expressing the above-described nucleic acid molecules. Within one embodiment, such vectors comprise a promoter operably linked to one of the above-described nucleic acid molecules. Within other embodiments, recombinant viral vectors are provided which are capable of directing the expression of one of the above described nucleic acid molecules. Representative examples of such viral vectors include retroviral vectors, adenoviral vectors, and herpes simplex virus vectors. Also provided

by the present invention are host cells containing one of the above-described recombinant vectors.

Within other aspects of the present invention, isolated Interleukin-1 Type 3 receptors are provided. Within one embodiment, such receptors have the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 562. Within another embodiment, the receptors have the sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 561. Within yet further aspects of the invention, isolated soluble Interleukin-1 Type 3 receptors are provided. Within one embodiment, the isolated soluble Interleukin-1 Type 3 receptors have the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 336. Within another embodiment, the soluble receptors have the sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 338.

Within other aspects of the invention, isolated antibodies capable of specifically binding to an Interleukin-1 Type 3 receptor are provided. Within one embodiment, the antibody may be selected from the group consisting of polyclonal antibodies, monoclonal antibodies, and antibody fragments. Within other embodiments, antibodies are provided which are capable of blocking the binding of IL-1 to an Interleukin-1 Type 3 receptor. Within preferred embodiments, the antibody is selected from the group consisting of murine and human antibodies. In addition to antibodies, the present invention also provides hybridomas which produces an antibody as described above.

Within yet another aspect of the present invention, nucleic acid molecules are provided which are capable of specifically hybridizing to a nucleic acid molecule encoding any of the Interleukin-1 Type 3 receptors described above. Such molecules may be between at least "y" nucleotides long, wherein "y" is any integer between 14 and 2044, and furthermore, may be selected suitable for use as probes or primers described below. Particularly preferred probes of the present invention are at least 18 nucleotides in length.

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings. In addition, various references are set forth below which describe in more detail certain procedures or compositions (e.g., plasmids, etc.), and are therefore incorporated by reference in their entirety.

### 35 Brief Description of the Drawings

Figure 1 schematically illustrates a rat IL-1 type 3 receptor.

Figure 2 is a table which lists the homology of a human IL-1 type 3 receptor with its rat homologue, and other interleukin receptors.

Figure 3 is a graph which shows stimulation of a reporter product via a human IL-1 type 3 receptor.

5 Figure 4 is a graph which shows the expression pattern of the IL-1 Type 3 receptor based upon RNA protection assays.

Figures 5A and B are two graphs which show inhibition of thymocyte proliferation by soluble IL-1 receptors.z

## 10 Detailed Description of the Invention

### Definitions

Prior to setting forth the invention, it may be helpful to an understanding thereof to set forth definitions of certain terms to be used hereinafter.

15 "Interleukin-1 Type 3 Receptors" ("IL-1-3R") refers to receptor proteins which bind Interleukin-1 ( $\alpha$  or  $\beta$ ), and, when expressed on a cell surface, transduce the signal provided by Interleukin-1 to the cell, thereby mediating a biological effect within the cell. In their native configuration, IL-1 Type 3 receptors exist as membrane bound proteins, consisting of an extracellular domain, transmembrane domain, and intracellular domain (see Figure 1). IL-1-3R may be distinguished from other Interleukin-1 receptors  
20 based upon criteria such as affinity of substrate binding, tissue distribution, and sequence homology. For example, IL-1-3R of the present invention should be greater than 50% homologous, preferably greater than 75% to 80% homologous, more preferably greater than 85% to 90% homologous, and most preferably greater than 92%, 95%, or 97% homologous to the IL-1-3R disclosed herein (e.g., Sequence I.D. No. 1). As utilized  
25 within the context of the present invention, IL1-3R should be understood to include not only the proteins which are disclosed herein, but substantially similar derivatives and analogs as discussed below.

"Soluble Interleukin-1 Type 3 Receptor" ("sIL1-3R") refers to a protein which has an amino acid sequence corresponding to the extracellular region of an  
30 Interleukin-1 Type 3 receptor. The extracellular region of IL-1-3R may be readily determined by a hydrophobicity analysis utilizing a computer program such as PROTEAN (DNASTAR, Madison, WI), or by an alignment analysis with other known type 1 and type 2 Interleukin-1 receptors.

35 "Nucleic acid molecule" refers to a nucleic acid polymer or nucleic acid sequence, which exists in the form of a separate fragment or as a component of a larger nucleic acid construct. The nucleic acid molecule must have been derived from nucleic acids isolated at least once in substantially pure form, (i.e., substantially free of



contaminating endogenous materials), and in a quantity or concentration enabling identification and recovery. Such sequences are preferably provided in the form of an open reading frame uninterrupted by internal nontranslated sequences, or introns. As utilized herein, nucleic acid molecules should be understood to include deoxyribonucleic acid ("DNA") molecules (including genomic and cDNA molecules), ribonucleic acid ("RNA") molecules, hybrid or chimeric nucleic acid molecules (e.g., DNA-RNA hybrids), and where appropriate, nucleic acid molecule analogs and derivatives (e.g., peptide nucleic acids ("PNA")). Nucleic acid molecules of the present invention may also comprise sequences of non-translated nucleic acids where such additional sequences do not interfere with manipulation or expression of the open reading frame (e.g., sequences which are 5' or 3' from the open reading frame).

"Recombinant expression vector" refers to a replicable nucleic acid construct used either to amplify or to express nucleic acid sequences which encode IL-1 Type 3, or sIL-1 Type 3 receptors. This construct comprises an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters, and (2) the structural or coding sequence of interest. The recombinant expression vector may also comprise appropriate transcription and translation initiation and termination sequences.

As noted above, the present invention provides isolated nucleic acid molecules encoding Interleukin-1 Type 3 receptors. One representative IL-1 Type 3 receptor which may be obtained utilizing the methods described herein (see, e.g., Example 1) is schematically illustrated in Figure 1. Briefly, this IL-1 Type 3 receptor (see Sequence I.D. Nos. 1 and 2) is composed of an Extracellular N-terminal Domain (amino acids 1 - 336), a Transmembrane Domain (amino acids 337 - 357), and a C-terminal Intracellular Domain (358 - 562).

Although the above IL-1 Type 3 receptor has been provided for purposes of illustration (see also Sequence I.D. Nos. 3 and 4), the present invention should not be so limited. In particular, "IL-1-3R" and "sIL-1-3R" as utilized herein should be understood to include a wide variety of IL-1 Type 3 receptors which are encoded by nucleic acid molecules that have substantial similarity to the sequences disclosed in Sequences I.D. Nos. 1 and 3. As utilized within the context of the present invention, nucleic acid molecules which encode IL-1 Type 3 receptors are deemed to be substantially similar to those disclosed herein if: (a) the nucleic acid sequence is derived from the coding region of a native IL-1 Type 3 receptor gene (including, for example, allelic variations of the sequences disclosed herein); (b) the nucleic acid sequence is capable of hybridization to nucleic acid sequences of the present invention under

conditions of either moderate (e.g., 50% formamide, 5 x SSPE, 5 x Denhardt's, 0.1% SDS, 100 ug/ml Salmon Sperm DNA, and a temperature of 42°C) or high stringency (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, NY, 1989); or (c) nucleic acid sequences are degenerate as a result of the genetic code to the nucleic acid sequences defined in (a) or (b). Furthermore, as noted above, although DNA molecules are primarily referred to herein, as should be evident to one of skill in the art given the disclosure provided herein, a wide variety of related nucleic acid molecules may also be utilized in various embodiments described herein, including for example, RNA, nucleic acid analogues, as well as chimeric nucleic acid molecules which may be composed of more than one type of nucleic acid.

In addition, as noted above, within the context of the present invention "IL-1 Type 3 receptors" and "soluble IL-1 Type 3 receptors" should be understood to include derivatives and analogs of the IL-1 Type 3 receptors described above. Such derivatives include allelic variants and genetically engineered variants that contain conservative amino acid substitutions and/or minor additions, substitutions or deletions of amino acids, the net effect of which does not substantially change the biological activity (e.g., signal transduction) or function of the IL-1 Type 3 receptor. Such derivatives are generally greater than about 50% homologous, preferably greater than 75% to 80% homologous, more preferably greater than 85% to 90% homologous, and most preferably greater than 92%, 95% or 97% homologous. Homology may be determined, for example, by comparing sequence information using the GAP computer program, version 6.0, available from the University of Wisconsin Genetics Computer Group (UWGCG).

The primary amino acid structure of IL-1 Type 3 receptors may also be modified by derivatizing amino acid side chains, and/or the amino or carboxy terminus with various functional groups, in order to allow for the formation of various conjugates (e.g., protein-IL-1-3R conjugates). Alternatively, conjugates of IL-1-3R (and sIL-1-3R) may be constructed by recombinantly producing fusion proteins. Such fusion proteins may comprise, for example, IL-1-3R-protein Z wherein protein Z is another cytokine receptor (e.g., IL-2R, IL-3R, IL-4R, IL-5R, IL-6R, IL-7R, IL-8R, IL-9R, IL-10R, IL-11R, IL-12R, IL-13R, IL-14R, IL-15R or TNF ( $\alpha$  or  $\beta$ ) receptor; see WO91/03553); a binding portion of an antibody; a toxin (as discussed below); or a protein or peptide which facilitates purification or identification of IL-1-3R (e.g., poly-His). For example, a fusion protein such as human IL-1-3R (His)<sub>n</sub> or sIL-1-3R (His)<sub>n</sub> may be constructed in order to allow purification of the protein via the poly-His residue, for example, on a NTA nickel-chelating column. The amino acid sequence of a IL-1 Type 3 receptor may



also be linked to the peptide Asp-Tyr-Lys-Asp-Asp-Asp-Lys (DYKDDDDK) (Sequence I.D. No. 5) (Hopp et al., *Bio/Technology* 6:1204, 1988) in order to facilitate purification of expressed recombinant protein.

The present invention also includes IL-1-3R (and sIL-1-3R) proteins which may be produced either with or without associated native-pattern glycosylation. For example, expression of IL-1-3R DNAs in bacteria such as *E. coli* provides non-glycosylated molecules. In contrast, IL-1-3R expressed in yeast or mammalian expression systems (as discussed below) may vary in both glycosylation pattern and molecular weight from native IL-1-3R, depending on the amino acid sequence and expression system which is utilized. In addition, functional mutants of mammalian IL-1-3R having inactivated glycosylation sites may also be produced in a homogeneous, reduced-carbohydrate form, utilizing oligonucleotide synthesis, site-directed mutagenesis, or random mutagenesis techniques. Briefly, N-glycosylation sites in eukaryotic proteins are generally characterized by the amino acid triplet Asn-A<sub>1</sub>-Z, where A<sub>1</sub> is any amino acid except Pro, and Z is Ser or Thr. In this triplet, asparagine provides a side chain amino group for covalent attachment of carbohydrate. Such sites may be eliminated by deleting Asn or Z, substituting another amino acid for Asn or for residue Z, or inserting a non-Z amino acid between A<sub>1</sub> and Z, or an amino acid other than Asn between Asn and A<sub>1</sub>.

Proteins which are substantially similar to IL-1-3R proteins may also be constructed by, for example, substituting or deleting various amino acid residues which are not required for biological activity. For example, cysteine residues may be deleted or replaced with other amino acids to prevent formation of incorrect intramolecular disulfide bridges upon renaturation. Similarly, adjacent dibasic amino acid residues may be modified for expression in yeast systems in which KEX2 protease activity is present.

Not all mutations in the nucleotide sequence which encodes IL-1-3R will be expressed in the final product. For example, nucleotide substitutions may be made in order to avoid secondary structure loops in the transcribed mRNA, or to provide codons that are more readily translated by the selected host, and thereby enhance expression within a selected host.

Generally, substitutions at the amino acid level should be made conservatively, i.e., the most preferred substitute amino acids are those which have characteristics resembling those of the residue to be replaced. When a substitution, deletion, or insertion strategy is adopted, the potential effect of the deletion or insertion on biological activity should be considered utilizing, for example, the signalling assay disclosed within the Examples.

Mutations which are made to the sequence of the nucleic acid molecules of the present invention should generally preserve the reading frame phase of the coding sequences. Furthermore, the mutations should preferably not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, which would adversely affect translation of the receptor mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation *per se* be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon, and the expressed IL-1-3R mutants screened for the biological activity. Representative methods for random mutagenesis include those described by Ladner et al. in U.S. Patent Nos. 5,096,815; 5,198,346; and 5,223,409.

As noted above, mutations may be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, site-directed mutagenesis procedures may be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik, *Bio Techniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); Sambrook et al. (*Molecular cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, 1989); and U.S. Patent Nos. 4,518,584 and 4,737,462, which are incorporated by reference herein.

IL-1 Type 3 receptors, as well as substantially similar derivatives or analogs may be used as therapeutic reagents, immunogens, reagents in receptor-based immunoassays, or as binding agents for affinity purification procedures. Moreover, IL-1 Type 3 receptors of the present invention may be utilized to screen compounds for IL-1 Type 3 receptor agonist or antagonistic activity. IL-1 Type 3 receptor proteins may also be covalently bound through reactive side groups to various insoluble substrates, such as cyanogen bromine-activated, bisoxirane-activated, carbonyldiimidazole-activated, or tosyl-activated, agarose structures, or by adsorbing to polyolefin surfaces (with or without glutaraldehyde cross-linking). Once bound to a substrate, IL-1-3R may be used to selectively bind (for purposes of assay or purification) anti-IL-1-3R antibodies or IL-

#### ISOLATION OF IL-1 TYPE 3 RECEPTOR CDNA CLONES

As noted above, the present invention provides isolated nucleic acid molecules which encode IL-1 Type 3 receptors. Briefly, nucleic acid molecules which encode IL-1 Type 3 receptors of the present invention may be readily isolated from a variety of warm-blooded animals, including for example, humans, macaques, horses, cattle, sheep, pigs, dogs, cats, rats and mice. Particularly preferred tissues from which nucleic acid molecules which encode IL-1 Type 3 receptors may be isolated include brain, kidney and lung. Nucleic acid molecules which encode IL-1 Type 3 receptors of the present invention may be readily isolated from conventionally prepared cDNA libraries (see, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, NY, 1989) or from commercially obtained libraries (e.g., Stratagene, LaJolla, Calif.) utilizing the disclosure provided herein. Particularly preferred methods for obtaining isolated DNA molecules which encode IL-1 Type 3 receptors of the present invention are described in more detail below in Example 1 (see also Sequence I.D. Nos. 1 and 3).

As noted above, within particularly preferred embodiments of the invention, isolated nucleic acid molecules are provided which encode human IL-1 Type 3 receptors. Briefly, such nucleic acid molecules may be readily obtained by probing a human cDNA library either with a specific sequence as described below in Example 1, or with a rat sequence (e.g., Sequence I.D. Nos. 2 or 4) under conditions of high stringency (e.g., 50% formamide, 5 x SSC, 5x Denharts, 0.1% SDS, 100 ug/ml salmon sperm DNA, at 42°C for 12 hours). This may be followed by extensive washing with 2x SSC containing 0.2% SDS at 50°C. Suitable cDNA libraries may be obtained from commercial sources (e.g., Stratagene, LaJolla, Calif.; or Clontech, Palo Alto, Calif., or prepared utilizing standard techniques (see, e.g., Sambrook et al., *supra*).

#### PRODUCTION OF RECOMBINANT IL-1 TYPE 3 RECEPTORS

As noted above, the present invention also provides recombinant expression vectors which include synthetic or cDNA-derived DNA fragments encoding IL-1 Type 3 receptors or substantially similar proteins, which are operably linked to suitable transcriptional or translation regulatory elements derived from mammalian, microbial, viral or insect genes. Such regulatory elements include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and, within preferred embodiments, sequences which control the termination of transcription and translation. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants may additionally be incorporated. DNA regions are

operably linked when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous and, in the case of secretory leaders, contiguous and in reading frame.

Expression vectors may also contain DNA sequences necessary to direct the secretion of a polypeptide of interest. Such DNA sequences may include at least one secretory signal sequence. Representative secretory signals include the alpha factor signal sequence (pre-pro sequence; Kurjan and Herskowitz, *Cell* 30:933-943, 1982; Kurjan et al., U.S. Patent No. 4,546,082; Brake, EP 116,201), the PHO5 signal sequence (Beck et al., WO 86/00637), the BAR1 secretory signal sequence (MacKay et al., U.S. Patent No. 4,613,572; MacKay, WO 87/002670), the SUC2 signal sequence (Carlson et al., *Mol. Cell. Biol.* 3:439-447, 1983), the  $\alpha$ -1-antitrypsin signal sequence (Kurachi et al., *Proc. Natl. Acad. Sci. USA* 78:6826-6830, 1981), the  $\beta$ -2 plasmin inhibitor signal sequence (Tone et al., *J. Biochem. (Tokyo)* 102:1033-1042, 1987), the tissue plasminogen activator signal sequence (Pennica et al., *Nature* 301:214-221, 1983), the *E. coli* PhoA signal sequence (Yuan et al., *J. Biol. Chem.* 265:13528-13552, 1990) or any of the bacterial signal sequences reviewed, for example, by Oliver (*Ann. Rev. Microbiol.* 39:615-649, 1985). Alternatively, a secretory signal sequence may be synthesized according to the rules established, for example, by von Heinje (*Eur. J. Biochem.* 133:17-21, 1983; *J. Mol. Biol.* 184:99-105, 1985; *Nuc. Acids Res.* 14:4683-4690, 1986).

For expression, a nucleic acid molecule encoding a IL-1 Type 3 receptor is inserted into a suitable expression vector, which in turn is used to transform or transfect appropriate host cells for expression. Host cells for use in practicing the present invention include mammalian, avian, plant, insect, bacterial and fungal cells. Preferred eukaryotic cells include cultured mammalian cell lines (e.g., rodent or human cell lines) and fungal cells, including species of yeast (e.g., *Saccharomyces* spp., particularly *S. cerevisiae*, *Schizosaccharomyces* spp., or *Kluyveromyces* spp.) or filamentous fungi (e.g., *Aspergillus* spp., *Neurospora* spp.). Strains of the yeast *Saccharomyces cerevisiae* are particularly preferred. Methods for producing recombinant proteins in a variety of prokaryotic and eukaryotic host cells are generally known in the art (see "Gene Expression Technology," *Methods in Enzymology*, Vol. 185, Goeddel (ed.), Academic Press, San Diego, Calif., 1990; see also, "Guide to Yeast Genetics and Molecular Biology," *Methods in Enzymology*, Guthrie and Fink (eds.)

Academic Press, San Diego, Calif., 1991). In general, a host cell will be selected on the basis of its ability to produce the protein of interest at a high level or its ability to carry out at least some of the processing steps necessary for the biological activity of the protein. In this way, the number of cloned DNA sequences which must be transfected  
5 into the host cell may be minimized and overall yield of biologically active protein may be maximized.

Suitable yeast vectors for use in the present invention include YRp7 (Struhl et al., *Proc. Natl. Acad. Sci. USA* 76:1035-1039, 1978), YEp13 (Broach et al., *Gene* 8:121-133, 1979), POT vectors (Kawasaki et al., U.S. Patent No. 4,931,373,  
10 which is incorporated by reference herein), pJDB249 and pJDB219 (Beggs, *Nature* 275:104-108, 1978) and derivatives thereof. Such vectors will generally include a selectable marker, which may be one of any number of genes that exhibit a dominant phenotype for which a phenotypic assay exists to enable transformants to be selected. Preferred selectable markers are those that complement host cell auxotrophy, provide  
15 antibiotic resistance or enable a cell to utilize specific carbon sources, and include *LEU2* (Broach et al., *ibid.*), *URA3* (Botstein et al., *Gene* 8:17, 1979), *HIS3* (Struhl et al., *ibid.*) or *POT1* (Kawasaki et al., *ibid.*). Another suitable selectable marker is the CAT gene, which confers chloramphenicol resistance on yeast cells.

Preferred promoters for use in yeast include promoters from yeast  
20 glycolytic genes (Hitzeman et al., *J. Biol. Chem.* 255:12073-12080, 1980; Alber and Kawasaki, *J. Mol. Appl. Genet.* 1:419-434, 1982; Kawasaki, U.S. Patent No. 4,599,311) or alcohol dehydrogenase genes (Young et al., in *Genetic Engineering of Microorganisms for Chemicals*, Hollaender et al. (eds.), p. 355, Plenum, New York, 1982; Ammerer, *Meth. Enzymol.* 101:192-201, 1983). In this regard, particularly  
25 preferred promoters are the *TPI1* promoter (Kawasaki, U.S. Patent No. 4,599,311, 1986) and the *ADH2-4<sup>c</sup>* promoter (Russell et al., *Nature* 304:652-654, 1983; Irani and Kilgore, U.S. Patent Application Serial No. 07/784,653, which is incorporated herein by reference). The expression units may also include a transcriptional terminator, such as the *TPI1* terminator (Alber and Kawasaki, *ibid.*).

30 In addition to yeast, proteins of the present invention can be expressed in filamentous fungi, for example, strains of the fungi *Aspergillus* (McKnight et al., U.S. Patent No. 4,935,349, which is incorporated herein by reference). Examples of useful promoters include those derived from *Aspergillus nidulans* glycolytic genes, such as the *ADH3* promoter (McKnight et al., *EMBO J.* 4:2093-2099, 1985) and the *tpiA* promoter.  
35 An example of a suitable terminator is the *ADH3* terminator (McKnight et al., *ibid.*, 1985). The expression units utilizing such components are cloned into vectors that are capable of insertion into the chromosomal DNA of *Aspergillus*.



Techniques for transforming fungi are well known in the literature, and have been described, for instance, by Beggs (*ibid.*), Hinnen et al. (*Proc. Natl. Acad. Sci. USA* 75:1929-1933, 1978), Yelton et al. (*Proc. Natl. Acad. Sci. USA* 81:1740-1747, 1984), and Russell (*Nature* 301:167-169, 1983). The genotype of the host cell will generally contain a genetic defect that is complemented by the selectable marker present on the expression vector. Choice of a particular host and selectable marker is well within the level of ordinary skill in the art. To optimize production of the heterologous proteins in yeast, for example, it is preferred that the host strain carries a mutation, such as the yeast *pep4* mutation (Jones, *Genetics* 85:23-33, 1977), which results in reduced proteolytic activity.

In addition to fungal cells, cultured mammalian cells may be used as host cells within the present invention. Preferred cultured mammalian cells for use in the present invention include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), BHK (ATCC No. CRL 1632), and 293 (ATCC No. CRL 1573; Graham et al., *J. Gen. Virol.* 36:59-72, 1977) cell lines. A preferred BHK cell line is the BHK 570 cell line (deposited with the American Type Culture Collection under accession number CRL 10314). In addition, a number of other mammalian cell lines may be used within the present invention, including Rat Hep I (ATCC No. CRL 1600), Rat Hep II (ATCC No. CRL 1548), TCMK (ATCC No. CCL 139), Human lung (ATCC No. CCL 75.1), Human hepatoma (ATCC No. HTB-52), Hep G2 (ATCC No. HB 8065), Mouse liver (ATCC No. CCL 29.1), NCTC 1469 (ATCC No. CCL 9.1), SP2/0-Ag14 (ATCC No. 1581), HIT-T15 (ATCC No. CRL 1777), Ltk<sup>-</sup> (ATCC) No. CCL 1.3) and RINm 5AHT<sub>2</sub>B (Orskov and Nielson, *FEBS* 229(1):175-178, 1988).

Mammalian expression vectors for use in carrying out the present invention should include a promoter capable of directing the transcription of a cloned gene or cDNA. Preferred promoters include viral promoters and cellular promoters. Viral promoters include the immediate early cytomegalovirus promoter (Boshart et al., *Cell* 41:521-530, 1985) and the SV40 promoter (Subramani et al., *Mol. Cell. Biol.* 1:854-864, 1981). Cellular promoters include the mouse metallothionein-1 promoter (Palmiter et al., U.S. Patent No. 4,579,821), a mouse V<sub>J</sub> promoter (Bergman et al., *Proc. Natl. Acad. Sci. USA* 81:7041-7045, 1983; Grant et al., *Nuc. Acids Res.* 15:5496, 1987) and a mouse V<sub>H</sub> promoter (Loh et al., *Cell* 33:85-93, 1983). A particularly preferred promoter is the major late promoter from Adenovirus 2 (Kaufman and Sharp, *Mol. Cell. Biol.* 2:1304-1319, 1982). Such expression vectors may also contain a set of RNA splice sites located downstream from the promoter and upstream from the DNA sequence encoding the peptide or protein of interest. Preferred RNA splice sites may be obtained from SV40, adenovirus and/or immunoglobulin genes. Alternatively, within



certain embodiments RNA splice sites may be located downstream from the DNA sequence encoding the peptide or protein of interest. Also contained in the expression vectors is a polyadenylation signal located downstream of the coding sequence of interest. Suitable polyadenylation signals include the early or late polyadenylation signals from SV40 (Kaufman and Sharp, *ibid.*), the polyadenylation signal from the Adenovirus 5 E1B region and the human growth hormone gene terminator (DeNoto et al., *Nuc. Acids Res.* 9:3719-3730, 1981). The expression vectors may include a noncoding viral leader sequence, such as the Adenovirus 2 tripartite leader, located between the promoter and the RNA splice sites. Preferred vectors may also include enhancer sequences, such as the SV40 enhancer and the mouse 1 enhancer (Gillies, *Cell* 33:717-728, 1983). Expression vectors may also include sequences encoding the adenovirus VA RNAs. Suitable vectors can be obtained from commercial sources (e.g., Invitrogen, San Diego, CA; Stratagene, La Jolla, CA).

Cloned DNA sequences may be introduced into cultured mammalian cells by, for example, calcium phosphate-mediated transfection (Wigler et al., *Cell* 14:725, 1978; Corsaro and Pearson, *Somatic Cell Genetics* 7:603, 1981; Graham and Van der Eb, *Virology* 52:456, 1973), electroporation (Neumann et al., *EMBO J.* 1:841-845, 1982), or DEAE-dextran mediated transfection (Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., NY, 1987), which are incorporated herein by reference. To identify cells that have stably integrated the cloned DNA, a selectable marker is generally introduced into the cells along with the gene or cDNA of interest. Preferred selectable markers for use in cultured mammalian cells include genes that confer resistance to drugs, such as neomycin, hygromycin, and methotrexate. The selectable marker may be an amplifiable selectable marker. Preferred amplifiable selectable markers are the DHFR gene and the neomycin resistance gene. Selectable markers are reviewed by Thilly (*Mammalian Cell Technology*, Butterworth Publishers, Stoneham, MA, which is incorporated herein by reference). The choice of selectable markers is well within the level of ordinary skill in the art.

Selectable markers may be introduced into the cell on a separate vector at the same time as the IL-1 Type 3 receptor sequence, or they may be introduced on the same vector. If on the same vector, the selectable marker and the IL-1 Type 3 receptor sequence may be under the control of different promoters or the same promoter, the latter arrangement producing a dicistronic message. Constructs of this type are known in the art (for example, Levinson and Simonsen, U.S. Patent No. 4,713,339). It may also be advantageous to add additional DNA, known as "carrier DNA" to the mixture which is introduced into the cells.

Transfected mammalian cells are allowed to grow for a period of time, typically 1-2 days, to begin expressing the DNA sequence(s) of interest. Drug selection is then applied to select for growth of cells that are expressing the selectable marker in a stable fashion. For cells that have been transfected with an amplifiable selectable marker the drug concentration may be increased in a stepwise manner to select for increased copy number of the cloned sequences, thereby increasing expression levels. Cells expressing the introduced sequences are selected and screened for production of the protein of interest in the desired form or at the desired level. Cells which satisfy these criteria may then be cloned and scaled up for production.

Preferred prokaryotic host cells for use in carrying out the present invention are strains of the bacteria *Escherichia coli*, although *Bacillus* and other genera are also useful. Techniques for transforming these hosts and expressing foreign DNA sequences cloned therein are well known in the art (see, e.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1982; or Sambrook et al., *supra*). Vectors used for expressing cloned DNA sequences in bacterial hosts will generally contain a selectable marker, such as a gene for antibiotic resistance, and a promoter that functions in the host cell. Appropriate promoters include the trp (Nichols and Yanofsky, *Meth. Enzymol.* 101:155-164, 1983), lac (Casadaban et al., *J. Bacteriol.* 143:971-980, 1980), and phage k (Queen, *J. Mol. Appl. Genet.* 2:1-10, 1983) promoter systems. Plasmids useful for transforming bacteria include pBR322 (Bolivar et al., *Gene* 2:95-113, 1977), the pUC plasmids (Messing, *Meth. Enzymol.* 101:20-78, 1983; Vieira and Messing, *Gene* 19:259-268, 1982), pCQV2 (Queen, *ibid.*), pMAL-2 (New England Biolabs, Beverly, MA) and derivatives thereof. Plasmids may contain both viral and bacterial elements.

Given the teachings provided herein, promoters, terminators and methods for introducing expression vectors encoding IL-1 Type 3 receptors of the present invention into plant, avian and insect cells would be evident to those of skill in the art. The use of baculoviruses, for example, as vectors for expressing heterologous DNA sequences in insect cells has been reviewed by Atkinson et al. (*Pestic. Sci.* 28:215-224, 1990). In addition, the use of *Agrobacterium rhizogenes* as vectors for expressing genes in plant cells has been reviewed by Sinkar et al. (*J. Biosci. (Bangalore)* 11:47-58, 1987).

Host cells containing DNA molecules of the present invention are then cultured to express a DNA molecule encoding a IL-1 Type 3 receptor. The cells are cultured according to standard methods in a culture medium containing nutrients required for growth of the chosen host cells. A variety of suitable media are known in the art and generally include a carbon source, a nitrogen source, essential amino acids,

vitamins and minerals, as well as other components, e.g., growth factors or serum, that may be required by the particular host cells. The growth medium will generally select for cells containing the DNA molecules by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker on the DNA construct or co-transfected with the DNA construct.

Suitable growth conditions for yeast cells, for example, include culturing in a chemically defined medium, comprising a nitrogen source, which may be a non-amino acid nitrogen source or a yeast extract, inorganic salts, vitamins and essential amino acid supplements at a temperature between 4°C and 37°C, with 30°C being particularly preferred. The pH of the medium is preferably maintained at a pH greater than 2 and less than 8, more preferably pH 5-6. Methods for maintaining a stable pH include buffering and constant pH control. Preferred agents for pH control include sodium hydroxide. Preferred buffering agents include succinic acid and Bis-Tris (Sigma Chemical Co., St. Louis, MO). Due to the tendency of yeast host cells to hyperglycosylate heterologous proteins, it may be preferable to express the IL-1 Type 3 receptors of the present invention in yeast cells having a defect in a gene required for asparagine-linked glycosylation. Such cells are preferably grown in a medium containing an osmotic stabilizer. A preferred osmotic stabilizer is sorbitol supplemented into the medium at a concentration between 0.1 M and 1.5 M, preferably at 0.5 M or 1.0 M. Cultured mammalian cells are generally cultured in commercially available serum-containing or serum-free media. Selection of a medium and growth conditions appropriate for the particular cell line used is within the level of ordinary skill in the art.

IL-1 Type 3 receptors may also be expressed in non-human transgenic animals, particularly transgenic warm-blooded animals. Methods for producing transgenic animals, including mice, rats, rabbits, sheep and pigs, are known in the art and are disclosed, for example, by Hammer et al. (*Nature* 315:680-683, 1985), Palmiter et al. (*Science* 222:809-814, 1983), Brinster et al. (*Proc. Natl. Acad. Sci. USA* 82:4438-4442, 1985), Palmiter and Brinster (*Cell* 41:343-345, 1985) and U.S. Patent No. 4,736,866, which are incorporated herein by reference. Briefly, an expression unit, including a DNA sequence to be expressed together with appropriately positioned expression control sequences, is introduced into pronuclei of fertilized eggs. Introduction of DNA is commonly done by microinjection. Integration of the injected DNA is detected by blot analysis of DNA from tissue samples, typically samples of tail tissue. It is generally preferred that the introduced DNA be incorporated into the germ line of the animal so that it is passed on to the animal's progeny.

Within particularly preferred embodiments of the invention, "knockout" animals may be developed from embryonic stem cells through the use of homologous

recombination (Capecchi, *Science* 244:1288-1292, 1989) or antisense oligonucleotide (Stein and Chen, *Science* 261(5124):1004-1012, 1993; Milligan et al., *Semin. Conc. Biol.* 3(6):391-398, 1992).

Within a preferred embodiment of the invention, a transgenic animal, such as a mouse, is developed by targeting a mutation to disrupt a IL-1 Type 3 receptor sequence (see Mansour et al., "Disruption of the proto-oncogene *int-2* in mouse embryo-derived stem cells: A general strategy for targeting mutations to non-selectable genes," *Nature* 336:348-352, 1988). Such animals may readily be utilized as a model to study the role of the IL-1 Type 3 receptor in metabolism.

10

#### SOLUBLE IL-1 TYPE 3 RECEPTORS AND RECEPTOR PEPTIDES

As noted above, the present invention also provides soluble IL-1 Type 3 receptors and receptor peptides. Within the context of the present invention, IL-1 Type 3 receptor peptides should be understood to include portions of a IL-1 Type 3 receptor or derivatives thereof discussed above, which do not contain transmembrane domains, and which are at least 8, and more preferably 10 or greater amino acids in length. Briefly, the structure of the IL-1 Type 3 receptor as well as putative transmembrane domains may be predicted from the primary translation products using the hydrophobicity plot function of, for example, PROTEAN (DNA STAR, Madison, WI), or according to the methods described by Kyte and Doolittle (*J. Mol. Biol.* 157:105-132, 1982). While not wishing to be bound by a graphical representation, based upon this hydrophobicity analysis, IL-1 Type 3 receptors are believed to have the general structure shown in Figure 1. In particular, these receptors are believed to comprise an extracellular amino-terminal domain, a transmembrane domain, and an intracellular domain.

Within one aspect of the invention, isolated IL-1 Type 3 receptor peptides are provided comprising the extracellular amino-terminal domain of a IL-1 Type 3 receptor. Within a preferred embodiment, an isolated IL-1 Type 3 receptor peptide is provided comprising the sequence of amino acids shown in Sequence I.D. No.2, from amino acid number 1 to amino acid number 336. Within other embodiments, isolated IL-1 Type 3 receptor peptides are provided comprising the sequence of amino acids shown in Sequence I.D. No. 4, from amino acid number 1 to amino acid number 338.

IL-1 Type 3 receptor peptides may be prepared by, among other methods, culturing suitable host/vector systems to produce the recombinant translation products of the present invention. Supernatants from such cell lines may then be treated by a variety of purification procedures in order to isolate the IL-1 Type 3 receptor

peptide. For example, the supernatant may be first concentrated using commercially available protein concentration filters, such as an Amicon or Millipore Pellicon ultrafiltration unit. Following concentration, the concentrate may be applied to a suitable purification matrix such as, for example, IL-1 or an anti-IL-1 Type 3 receptor antibody bound to a suitable support. Alternatively, anion or cation exchange resins may be employed in order to purify the receptor or peptide. Finally, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps may be employed to further purify the IL-1 Type 3 receptor peptide.

Alternatively, IL-1 Type 3 receptor peptides may also be prepared utilizing standard polypeptide synthesis protocols, and purified utilizing the above-described procedures.

A IL-1 Type 3 receptor peptide is deemed to be "isolated" or purified within the context of the present invention, if only a single band is detected subsequent to SDS-polyacrylamide gel analysis followed by staining with Coomassie Brilliant Blue.

#### ANTIBODIES TO IL-1 TYPE 3 RECEPTORS

Within one aspect of the present invention, IL-1 Type 3 receptors, including derivatives thereof, as well as portions or fragments of these proteins such as the IL-1 Type 3 receptor peptides discussed above, may be utilized to prepare antibodies which specifically bind to IL-1 Type 3 receptors. Within the context of the present invention the term "antibodies" includes polyclonal antibodies, monoclonal antibodies, fragments thereof such as F(ab')<sub>2</sub> and Fab fragments, as well as recombinantly produced binding partners. These binding partners incorporate the variable regions from a gene which encodes a specifically binding monoclonal antibody. Antibodies are defined to be specifically binding if they bind to the IL-1 Type 3 receptor with a  $K_A$  of greater than or equal to  $10^7 \text{ M}^{-1}$  and preferably greater than or equal to  $10^8 \text{ M}^{-1}$ , and bind to IL-1 Type I or Type II receptors with an affinity of less than  $K_A 10^7 \text{ M}^{-1}$ , and preferably less than  $10^5 \text{ M}^{-1}$  or  $10^3 \text{ M}^{-1}$ . The affinity of a monoclonal antibody or binding partner may be readily determined by one of ordinary skill in the art (see Scatchard, *Ann. N.Y. Acad. Sci.* 51:660-672, 1949).

Polyclonal antibodies may be readily generated by one of ordinary skill in the art from a variety of warm-blooded animals such as horses, cows, goats, sheep, dogs, chickens, rabbits, mice, or rats. Briefly, the IL-1 Type 3 receptor is utilized to immunize the animal through intraperitoneal, intramuscular, intraocular, or subcutaneous injections. The immunogenicity of a IL-1 Type 3 receptor or IL-1 Type 3 receptor peptide may be increased through the use of an adjuvant such as Freund's complete or incomplete adjuvant. Following several booster immunizations, small samples of serum



are collected and tested for reactivity to the IL-1 Type 3 receptor. A variety of assays may be utilized in order to detect antibodies which specifically bind to a IL-1 Type 3 receptor. Exemplary assays are described in detail in *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988. Representative  
5 examples of such assays include: Countercurrent Immuno-Electrophoresis (CIEP), Radioimmunoassays, Radioimmunoprecipitations, Enzyme-Linked Immuno-Sorbent Assays (ELISA), Dot Blot assays, Inhibition or Competition assays, and sandwich assays (see U.S. Patent Nos. 4,376,110 and 4,486,530; see also *Antibodies: A Laboratory Manual, supra*). Particularly preferred polyclonal antisera will give a signal  
10 that is at least three times greater than background. Once the titer of the animal has reached a plateau in terms of its reactivity to the IL-1 Type 3 receptor, larger quantities of polyclonal antisera may be readily obtained either by weekly bleedings, or by exsanguinating the animal.

Monoclonal antibodies may also be readily generated using well-known  
15 techniques (see U.S. Patent Nos. RE 32,011, 4,902,614, 4,543,439, and 4,411,993; see also *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, Kennett, McKearn, and Bechtol (eds.), 1980, and *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988). Briefly, within one embodiment a subject animal such as a rat or mouse is  
20 injected with a form of IL-1 Type 3 receptor suitable for generating an immune response against the IL-1 Type 3 receptor. Representative examples of suitable forms include, among others, cells which express the IL-1 Type 3 receptor, or peptides which are based upon the IL-1 Type 3 receptor sequence. Additionally, many techniques are known in the art for increasing the resultant immune response, for example, by coupling the  
25 receptor or receptor peptides to another protein such as ovalbumin or keyhole limpet hemocyanin (KLH), or through the use of adjuvants such as Freund's complete or incomplete adjuvant. The initial immunization may be through intraperitoneal, intramuscular, intraocular, or subcutaneous routes.

Between one and three weeks after the initial immunization the animal  
30 may be reimmunized with another booster immunization. The animal may then be test bled and the serum tested for binding to the IL-1 Type 3 receptor using assays as described above. Additional immunizations may also be accomplished until the animal has plateaued in its reactivity to the IL-1 Type 3 receptor. The animal may then be given a final boost of IL-1 Type 3 receptor or IL-1 Type 3 receptor peptide, and three  
35 to four days later sacrificed. At this time, the spleen and lymph nodes may be harvested and disrupted into a single cell suspension by passing the organs through a mesh screen or by rupturing the spleen or lymph node membranes which encapsidate the cells.



Within one embodiment the red cells are subsequently lysed by the addition of a hypotonic solution, followed by immediate return to isotonicity.

Within another embodiment, suitable cells for preparing monoclonal antibodies are obtained through the use of *in vitro* immunization techniques. Briefly, an animal is sacrificed, and the spleen and lymph node cells are removed as described above. A single cell suspension is prepared, and the cells are placed into a culture containing a form of the IL-1 Type 3 receptor that is suitable for generating an immune response as described above. Subsequently, the lymphocytes are harvested and fused as described below.

Cells which are obtained through the use of *in vitro* immunization or from an immunized animal as described above may be immortalized by transfection with a virus such as the Epstein-Barr virus (EBV) (see Glasky and Reading, *Hybridoma* 8(4):377-389, 1989). Alternatively, within a preferred embodiment, the harvested spleen and/or lymph node cell suspensions are fused with a suitable myeloma cell in order to create a "hybridoma" which secretes monoclonal antibodies. Suitable myeloma lines are preferably defective in the construction or expression of antibodies, and are additionally syngeneic with the cells from the immunized animal. Many such myeloma cell lines are well known in the art and may be obtained from sources such as the American Type Culture Collection (ATCC), Rockville, Maryland (see *Catalogue of Cell Lines & Hybridomas*, 6th ed., ATCC, 1988). Representative myeloma lines include: for humans, UC 729-6 (ATCC No. CRL 8061), MC/CAR-Z2 (ATCC No. CRL 8147), and SKO-007 (ATCC No. CRL 8033); for mice, SP2/0-Ag14 (ATCC No. CRL 1581), and P3X63Ag8 (ATCC No. TIB 9); and for rats, Y3-Ag1.2.3 (ATCC No. CRL 1631), and YB2/0 (ATCC No. CRL 1662). Particularly preferred fusion lines include NS-1 (ATCC No. TIB 18) and P3X63 - Ag 8.653 (ATCC No. CRL 1580), which may be utilized for fusions with either mouse, rat, or human cell lines. Fusion between the myeloma cell line and the cells from the immunized animal may be accomplished by a variety of methods, including the use of polyethylene glycol (PEG) (see *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988) or electrofusion (see Zimmerman and Vienken, *J. Membrane Biol.* 67:165-182, 1982).

Following the fusion, the cells are placed into culture plates containing a suitable medium, such as RPMI 1640 or DMEM (Dulbecco's Modified Eagles Medium) (JRH Biosciences, Lenexa, KS). The medium may also contain additional ingredients, such as Fetal Bovine Serum ("FBS," *i.e.*, from Hyclone, Logan, Utah, or JRH Biosciences), thymocytes which were harvested from a baby animal of the same species as was used for immunization, or agar to solidify the medium. Additionally, the medium should contain a reagent which selectively allows for the growth of fused spleen and

myeloma cells. Particularly preferred is the use of HAT (hypoxanthine, aminopterin, and thymidine) (Sigma Chemical Co., St. Louis, MO). After about seven days, the resulting fused cells or hybridomas may be screened in order to determine the presence of antibodies which recognize the IL-1 Type 3 receptor. Following several clonal dilutions and reassays, a hybridoma producing antibodies which bind to IL-1 Type 3 receptor may be isolated.

Other techniques may also be utilized to construct monoclonal antibodies (see Huse et al., "Generation of a Large Combinational Library of the Immunoglobulin Repertoire in Phage Lambda," *Science* 246:1275-1281, December 1989; see also Sastry et al., "Cloning of the Immunological Repertoire in *Escherichia coli* for Generation of Monoclonal Catalytic Antibodies: Construction of a Heavy Chain Variable Region-Specific cDNA Library," *Proc. Natl. Acad. Sci. USA* 86:5728-5732, August 1989; see also Altling-Mees et al., "Monoclonal Antibody Expression Libraries: A Rapid Alternative to Hybridomas," *Strategies in Molecular Biology* 3:1-9, January 1990; these references describe a commercial system available from Stratacyte, La Jolla, California, which enables the production of antibodies through recombinant techniques). Briefly, mRNA is isolated from a B cell population and utilized to create heavy and light chain immunoglobulin cDNA expression libraries in the kIMMUNOZAP(H) and kIMMUNOZAP(L) vectors. These vectors may be screened individually or co-expressed to form Fab fragments or antibodies (see Huse et al., *supra*; see also Sastry et al., *supra*). Positive plaques may subsequently be converted to a non-lytic plasmid which allows high level expression of monoclonal antibody fragments from *E. coli*.

Similarly, binding partners may also be constructed utilizing recombinant DNA techniques to incorporate the variable regions of a gene which encodes a specifically binding antibody. The construction of these proteins may be readily accomplished by one of ordinary skill in the art (see Larrick et al., "Polymerase Chain Reaction Using Mixed Primers: Cloning of Human Monoclonal Antibody Variable Region Genes From Single Hybridoma Cells," *Biotechnology* 7:934-938, September 1989; Riechmann et al., "Reshaping Human Antibodies for Therapy," *Nature* 332:323-327, 1988; Roberts et al., "Generation of an Antibody with Enhanced Affinity and Specificity for its Antigen by Protein Engineering," *Nature* 328:731-734, 1987; Verhoeyen et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity," *Science* 239:1534-1536, 1988; Chaudhary et al., "A Recombinant Immunotoxin Consisting of Two Antibody Variable Domains Fused to *Pseudomonas* Exotoxin," *Nature* 339:394-397, 1989; see also, U.S. Patent No. 5,132,405 entitled "Biosynthetic Antibody Binding Sites"), given the disclosure provided herein. Briefly, within one embodiment, DNA molecules encoding IL-1 Type 3 receptor-specific antigen binding

domains are amplified from hybridomas which produce a specifically binding monoclonal antibody, and inserted directly into the genome of a cell which produces human antibodies (*see* Verhoeyen et al., *supra*; *see also* Reichmann et al., *supra*). This technique allows the antigen-binding site of a specifically binding mouse or rat monoclonal antibody to be transferred into a human antibody. Such antibodies are preferable for therapeutic use in humans because they are not as antigenic as rat or mouse antibodies.

Alternatively, the antigen-binding sites (variable region) may be either linked to, or inserted into, another completely different protein (*see* Chaudhary et al., *supra*), resulting in a new protein with antigen-binding sites of the antibody as well as the functional activity of the completely different protein. As one of ordinary skill in the art will recognize, the antigen-binding sites or IL-1 Type 3 receptor binding domain of the antibody may be found in the variable region of the antibody. Furthermore, DNA sequences which encode smaller portions of the antibody or variable regions which specifically bind to mammalian IL-1 Type 3 receptor may also be utilized within the context of the present invention. These portions may be readily tested for binding specificity to the IL-1 Type 3 receptor utilizing assays described below.

Within a preferred embodiment, genes which encode the variable region from a hybridoma producing a monoclonal antibody of interest are amplified using oligonucleotide primers for the variable region. These primers may be synthesized by one of ordinary skill in the art, or may be purchased from commercially available sources. Stratacyte (La Jolla, CA) sells primers for mouse and human variable regions including, among others, primers for  $V_{H\alpha}$ ,  $V_{H\beta}$ ,  $V_{H\gamma}$ ,  $V_{H\delta}$ ,  $C_{H1}$ ,  $V_L$  and  $C_L$  regions. These primers may be utilized to amplify heavy or light chain variable regions, which may then be inserted into vectors such as IMMUNOZAP\*(H) or IMMUNOZAP\*(L) (Stratacyte), respectively. These vectors may then be introduced into *E. coli* for expression. Utilizing these techniques, large amounts of a single-chain protein containing a fusion of the  $V_H$  and  $V_L$  domains may be produced (*see* Bird et al., *Science* 242:423-426, 1988).

Other "antibodies" which may also be prepared utilizing the disclosure provided herein, and thus which are also deemed to fall within the scope of the present invention include humanized antibodies (*e.g.*, U.S. Patent No. 4,816,567 and WO 94/10332), micobodies (*e.g.*, WO 94/09817) and transgenic antibodies (*e.g.*, GB 2 272 440).

Once suitable antibodies have been obtained, they may be isolated or purified by many techniques well known to those of ordinary skill in the art (*see Antibodies: A Laboratory Manual, supra*). Suitable techniques include peptide or

protein affinity columns, HPLC or RP-HPLC, purification on protein A or protein G columns, or any combination of these techniques. Within the context of the present invention, the term "isolated" as used to define antibodies or binding partners means "substantially free of other blood components."

5           Antibodies of the present invention have many uses. For example, antibodies may be utilized in flow cytometry to sort IL-1 Type 3 receptor-bearing cells, or to histochemically stain IL-1 Type 3 receptor-bearing tissues. Briefly, in order to detect IL-1 Type 3 receptors on cells, the cells (or tissue) are incubated with a labeled antibody which specifically binds to IL-1 Type 3 receptors, followed by detection of the  
10   presence of bound antibody. These steps may also be accomplished with additional steps such as washings to remove unbound antibody. Representative examples of suitable labels, as well as methods for conjugating or coupling antibodies to such labels are described in more detail below.

          In addition, purified antibodies may also be utilized therapeutically to  
15   block the binding of IL-1 or other IL-1 Type 3 receptor substrates to the IL-1 Type 3 receptor *in vitro* or *in vivo*. As noted above, a variety of assays may be utilized to detect antibodies which block or inhibit the binding of IL-1 to the IL-1 Type 3 receptor, including *inter alia*, inhibition and competition assays noted above. Within one embodiment, monoclonal antibodies (prepared as described above) are assayed for  
20   binding to the IL-1 Type 3 receptor in the absence of IL-1, as well as in the presence of varying concentrations of IL-1. Blocking antibodies are identified as those which, for example, bind to IL-1 Type 3 receptors and, in the presence of IL-1, block or inhibit the binding of IL-1 to the IL-1 Type 3 receptor.

          Antibodies of the present invention may also be coupled or conjugated to  
25   a variety of other compounds (or labels) for either diagnostic or therapeutic use. Such compounds include, for example, toxic molecules, molecules which are nontoxic but which become toxic upon exposure to a second compound, and radionuclides. Representative examples of such molecules are described in more detail below.

          Antibodies which are to be utilized therapeutically are preferably  
30   provided in a therapeutic composition comprising the antibody or binding partner and a physiologically acceptable carrier or diluent. Suitable carriers or diluents include, among others, neutral buffered saline or saline, and may also include additional excipients or stabilizers such as buffers, sugars such as glucose, sucrose, or dextrose, chelating agents such as EDTA, and various preservatives.

### LABELS

The nucleic acid molecules, antibodies, and IL-1 Type 3 receptors (including sIL-1 3R) of the present invention may be labeled or conjugated (either through covalent or non-covalent means) to a variety of labels or other molecules, including for example, fluorescent markers, enzyme markers, toxic molecules, molecules which are nontoxic but which become toxic upon exposure to a second compound, and radionuclides.

Representative examples of fluorescent labels suitable for use within the present invention include, for example, Fluorescein Isothiocyanate (FITC), Rhodamine, Texas Red, Luciferase and Phycoerythrin (PE). Particularly preferred for use in flow cytometry is FITC which may be conjugated to purified antibody according to the method of Keltkamp in "Conjugation of Fluorescein Isothiocyanate to Antibodies. I. Experiments on the Conditions of Conjugation," *Immunology* 18:865-873, 1970. (See also Keltkamp, "Conjugation of Fluorescein Isothiocyanate to Antibodies. II. A Reproducible Method," *Immunology* 18:875-881, 1970; and Goding, "Conjugation of Antibodies with Fluorochromes: Modification to the Standard Methods," *J. Immunol. Methods* 13:215-226, 1970.) For histochemical staining, HRP, which is preferred, may be conjugated to the purified antibody according to the method of Nakane and Kawaoi ("Peroxidase-Labeled Antibody: A New Method of Conjugation," *J. Histochem. Cytochem.* 22:1084-1091, 1974; see also, Tijssen and Kurstak, "Highly Efficient and Simple Methods for Preparation of Peroxidase and Active Peroxidase Antibody Conjugates for Enzyme Immunoassays," *Anal. Biochem.* 136:451-457, 1984).

Representative examples of enzyme markers or labels include alkaline phosphatase, horse radish peroxidase, and  $\beta$ -galactosidase. Representative examples of toxic molecules include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, pokeweed antiviral protein, tritin, Shigella toxin, and Pseudomonas exotoxin A. Representative examples of molecules which are nontoxic, but which become toxic upon exposure to a second compound include thymidine kinases such as HSVTK and VZVTk. Representative examples of radionuclides include Cu-64, Ga-67, Ga-68, Zr-89, Ru-97, Tc-99m, Rh-105, Pd-109, In-111, I-123, I-125, I-131, Re-186, Re-188, Au-198, Au-199, Pb-203, At-211, Pb-212 and Bi-212.

As will be evident to one of skill in the art given the disclosure provided herein, the above described nucleic acid molecules, antibodies, and IL-1 Type 3 receptors may also be labeled with other molecules such as colloidal gold, as well either member of a high affinity binding pair (e.g., avidin-biotin).



DIAGNOSTIC USE OF IL-1 TYPE 3 RECEPTOR SEQUENCES

Within another aspect of the present invention, probes and primers are provided for detecting IL-1 Type 3 receptors. Within one embodiment of the invention, probes are provided which are capable of hybridizing to IL-1 Type 3 receptor DNA or RNA. For purposes of the present invention, probes are "capable of hybridizing" to IL-1 Type 3 receptor DNA if they hybridize to Sequence I.D. Nos. 1 or 3 under conditions of moderate or high stringency (*see* Sambrook et al., *supra*); but not to IL-1 Type I or Type II receptor nucleic acid sequences. Preferably, the probe may be utilized to hybridize to suitable nucleotide sequences in the presence of 50% formamide, 5x SSPE, 5x Denhardt's, 0.1% SDS and 100 ug/ml Salmon Sperm DNA at 42°C, followed by a first wash with 2x SSC at 42°C, and a second wash with 0.2x SSC at 55 to 60°C.

Probes of the present invention may be composed of either deoxyribonucleic acids (DNA) ribonucleic acids (RNA), nucleic acid analogues, or any combination of these, and may be as few as about 12 nucleotides in length, usually about 14 to 18 nucleotides in length, and possibly as large as the entire sequence of the IL-1 Type 3 receptor. Selection of probe size is somewhat dependent upon the use of the probe. For example, in order to determine the presence of various polymorphic forms of the IL-1 Type 3 receptor within an individual, a probe comprising virtually the entire length of the IL-1 Type 3 receptor coding sequence is preferred. IL-1 Type 3 receptor probes may be utilized to identify polymorphisms linked to the IL-1 Type 3 receptor gene (*see, for example*, Weber, *Genomics* 7:524-530, 1990; and Weber and May, *Amer. J. Hum. Gen.* 44:388-396, 1989). Such polymorphisms may be associated with inherited diseases such as diabetes.

Probes may be constructed and labeled using techniques which are well known in the art. Shorter probes of, for example, 12 or 14 bases may be generated synthetically. Longer probes of about 75 bases to less than 1.5 kb are preferably generated by, for example, PCR amplification in the presence of labeled precursors such as <sup>32</sup>P-dCTP, digoxigenin-dUTP, or biotin-dATP. Probes of more than 1.5 kb are generally most easily amplified by transfecting a cell with a plasmid containing the relevant probe, growing the transfected cell into large quantities, and purifying the relevant sequence from the transfected cells (*see* Sambrook et al., *supra*).

Probes may be labeled by a variety of markers, including, for example, radioactive markers, fluorescent markers, enzymatic markers, and chromogenic markers. The use of <sup>32</sup>P is particularly preferred for marking or labeling a particular probe.

Probes of the present invention may also be utilized to detect the presence of a IL-1 Type 3 receptor mRNA or DNA within a sample. However, if IL-1 Type 3 receptors are present in only a limited number, or if it is desired to detect a



selected mutant sequence which is present in only a limited number, or if it is desired to clone a IL-1 Type 3 receptor from a selected warm-blooded animal, then it may be beneficial to amplify the relevant sequence such that it may be more readily detected or obtained.

5 A variety of methods may be utilized in order to amplify a selected sequence, including, for example, RNA amplification (*see* Lizardi et al., *Bio/Technology* 6:1197-1202, 1988; Kramer et al., *Nature* 339:401-402, 1989; Lomeli et al., *Clinical Chem.* 35(9):1826-1831, 1989; U.S. Patent No. 4,786,600), and DNA amplification utilizing Polymerase Chain Reaction ("PCR") (*see* U.S. Patent Nos. 4,683,195,  
10 4,683,202, and 4,800,159) (*see also*, U.S. Patent Nos. 4,876,187, and 5,011,769, which describe an alternative detection/amplification system comprising the use of scissile linkages).

Within a particularly preferred embodiment, PCR amplification is utilized to detect or obtain a IL-1 Type 3 receptor DNA. Briefly, as described in greater detail  
15 below, a DNA sample is denatured at 95°C in order to generate single stranded DNA. Specific primers, as discussed below, are then annealed at 37°C to 70°C, depending on the proportion of AT/GC in the primers. The primers are extended at 72°C with Taq polymerase in order to generate the opposite strand to the template. These steps constitute one cycle, which may be repeated in order to amplify the selected sequence.

20 Primers for the amplification of a selected sequence should be selected from sequences which are highly specific and form stable duplexes with the target sequence. The primers should also be non-complementary, especially at the 3' end, should not form dimers with themselves or other primers, and should not form secondary structures or duplexes with other regions of DNA. In general, primers of  
25 about 18 to 20 nucleotides are preferred, and may be easily synthesized using techniques well known in the art.

#### PHARMACEUTICAL COMPOSITIONS AND THERAPEUTIC USES

As noted above, the present invention provides pharmaceutical  
30 compositions, as well as methods for using the same (for either prophylactic or therapeutic use). Briefly, the pharmaceutical compositions of the present invention may comprise an IL-1 3R, sIL-1 3R, antibody which is capable of specifically binding IL-1 3R, IL-1 3R antagonists or agonists, in combination with a pharmaceutically acceptable carrier, diluent, or excipient. Such compositions may comprise buffers such  
35 as neutral buffered saline, phosphate buffered saline and the like, carbohydrates such as glucose, mannose, sucrose or dextrose, proteins, polypeptides or amino acids, antioxidants, chelating agents such as EDTA or glutathione, and preservatives.

Compositions of the present invention may be formulated for the manner of administration indicated, including for example, for oral, nasal, venous, vaginal or rectal administration. Within other embodiments, the compositions may be administered as part of a sustained release implant (e.g., intra-articularly). Within yet other  
5   embodiments, the compositions may be formulated as a lyophilizate, utilizing appropriate excipients which provide stability as a lyophilizate, and subsequent to rehydration.

Pharmaceutical compositions of the present invention may be utilized in order to treat a wide variety of diseases including, for example, immune-associated diseases such as rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis,  
10   myasthenia gravis, scleritis, scleroderma, septic shock, allograft rejection, and graft versus host (GVH) disease. In particular, pharmaceutical compositions of the present invention may be administered in a manner appropriate to the disease to be treated (or prevented). Although appropriate dosages may be determined by clinical trials, the quantity and frequency of administration will be determined by such factors as the  
15   condition of the patient, and the type and severity of the patient's disease.

Within other aspects of the present invention, viral vectors are provided which may be utilized to treat diseases wherein either the IL-1 Type 3 receptor (or a mutant IL-1 Type 3 receptor) is over-expressed, or where no IL-1 Type 3 receptor is expressed. Briefly, within one embodiment of the invention, viral vectors are provided  
20   which direct the production of antisense IL-1 Type 3 receptor RNA, in order to prohibit the over-expression of IL-1 Type 3 receptors, or the expression of mutant IL-1 Type 3 receptors. Within another embodiment, viral vectors are provided which direct the expression of IL-1 Type 3 receptor cDNA. Viral vectors suitable for use in the present invention include, among others, recombinant vaccinia vectors (U.S. Patent Nos.  
25   4,603,112 and 4,769,330), recombinant pox virus vectors (PCT Publication No. WO 89/01973), and preferably, recombinant retroviral vectors ("Recombinant Retroviruses with Amphotropic and Ecotropic Host Ranges," PCT Publication No. WO 90/02806; "Retroviral Packaging Cell Lines and Processes of Using Same," PCT Publication No. WO 89/07150; and "Antisense RNA for Treatment of Retroviral Disease States," PCT  
30   Publication No. WO/03451), and herpesvirus vectors (Kit, *Adv. Exp. Med. Biol.* 215:219-236, 1989; U.S. Patent No. 5,288,641).

Within various embodiments of the invention, the above-described compositions may be administered *in vivo*, or *ex vivo*. Representative routes for *in vivo* administration include intradermally ("i.d."), intracranially ("i.c."), intraperitoneally  
35   ("i.p."), intrathecally ("i.t."), intravenously ("i.v."), subcutaneously ("s.c.") or intramuscularly ("i.m.).

Within other embodiments of the invention, the vectors which contain or express nucleic acid molecules of the present invention, or even the nucleic acid molecules themselves, may be administered by a variety of alternative techniques, including for example direct DNA injection (Acsadi et al., *Nature* 352:815-818, 1991);  
5 microprojectile bombardment (Williams et al., *PNAS* 88:2726-2730, 1991); liposomes (Pickering et al., *Circ.* 89(1):13-21, 1994; and Wang et al., *PNAS* 84:7851-7855, 1987); lipofection (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1989); DNA ligand (Wu et al., *J. of Biol. Chem.* 264:16985-16987, 1989); administration of DNA linked to killed adenovirus (Michael et al., *J. Biol. Chem.* 268(10):6866-6869, 1993; and  
10 Curiel et al., *Hum. Gene Ther.* 3(2):147-154, 1992), retrotransposons, cytofectin-mediated introduction (DMRIE-DOPE, Vical, Calif.) and transferrin-DNA complexes (Zenke).

The following examples are offered by way of illustration, and not by  
15 way of limitation.

EXAMPLESEXAMPLE 1

## ISOLATION OF INTERLEUKIN-1 TYPE 3 RECEPTOR cDNA

5

A. Isolation of Interleukin-1 Type 3 Receptor cDNA From a RatLung cDNA Library

Male Sprague-Dawley rats (Madison, WI) weighing between 175-250 gm are decapitated, and the lungs excised. Total RNA is then isolated from the lung utilizing a Promega RNAgents Total RNA Kit (catalog #Z5110, Promega, Wisc.) according to the manufacturers instructions, followed by the isolation of poly A+ RNA utilizing a Promega PolyAtract kit (catalog # Z5420). A cDNA phage library is then prepared utilizing a Giga-Pack Gold library construction kit according to the manufacturers' instructions (catalog #237611, Stratagene, LaJolla, Calif.), which is in turn plated and screened essentially as described by Sambrook et al., (Molecular Cloning) with oligonucleotide (5'-CTTCAACTGC ACATACCCTC CAGTAACAAA CGGGGCAGTG AATCTGACAT-3') (Sequence I.D. No. 6). This oligonucleotide is complementary to nucleotides 211-260 of the rat IL-1 Type 3 receptor cDNA sequence shown in Sequence I.D. No. 3.

The phage library is rescreened until a single pure phage isolate is obtained. The phage is then grown on bacterial host XL1-Blue (Stratagene, LaJolla, Calif.), and plasmid DNA is excised with ExAssist helper phage (Stratagene) in SOLR cells. The SOLR cells are then plated, and plasmid DNA is isolated and sequenced utilizing the Sanger dideoxy protocol.

A rat IL-1 Type 3 receptor cDNA sequence that may be obtained utilizing this procedure is set forth in Sequence I.D. No. 3.

B. Isolation of Interleukin-1 Type 3 Receptor cDNA From a30 Commerically Available Rat cDNA Library

IL-1 Type 3 receptor cDNA can also be isolated from commercially available rat cDNA libraries. For example, two million plaques from a rat phage library (Clontech, catalog #RL1048a) may be plated according to the manufacturer's instructions, and screened with oligonucleotide Sequence I.D. No. 6 essentially as described above.

A rat IL-1 Type 3 receptor cDNA sequence that may be obtained utilizing this procedure is set forth in Sequence I.D. No.3.

C. Isolation of IL-1 Type 3 Receptor cDNA From a Human cDNA Library

IL-1 Type 3 receptor cDNA can also be isolated from commercially available human cDNA libraries. Briefly, approximately two million plaques from a human phage library (Clontech, catalog #HL1158a) are plated according to the manufacturers instructions, and screened with oligonucleotide (5'-CCTCCCATAA CATCTGGGGA AGTCAGTGTA ACATGGTATA AAAATTCTAG C-3') (Sequence I.D. No. 7) essentially as described above. This oligonucleotide is complementary to nucleotides 260-310 of the human IL-1 Type 3 receptor cDNA sequence shown in Sequence I.D. No. 1.

The phage library is rescreened and isolated as described above. The human sequence that is obtained utilizing this procedure is approximately 89.1% identical at the nucleotide level and 89.2% identical at the amino acid level to that of the common region of the above-described rat IL-1 Type 3 receptors.

EXAMPLE 2

EXPRESSION OF IL-1 TYPE 3 RECEPTOR CDNA

A. Expression of Rat Interleukin-1 Type 3 Receptor

In order to express IL-1 Type 3 receptor cDNA, a mammalian cell expression vector (pCDM7amp) is first constructed. Briefly, pCDM7amp is a DNA plasmid which contains 1) an ampicillin resistance gene that provides for selection in prokaryotic cells, 2) a bacterial origin of replication which allows propagation and amplification in host bacterial cells, 3) a CMV (cytomegalovirus) promoter which sponsors transcription in mammalian cells, 4) a multiple cloning site (MCS), which is a series of adjacent restriction sites in the DNA sequence that are useful for the insertion of appropriate DNA fragments, and 5) a SV 40 T-antigen splice and polyadenylation site.

pCDM7-Amp is constructed from pCDM8 (Seed, Nature 329:840-842, 1987; Seed and Aruffo, Proc. Natl. Acad. Sci. 84:3365-3369, 1987; Thomsen et al., Cell 63:485-493, 1990; Bernot and Auffray, Proc. Natl. Acad. Sci 88:2550-2554, 1991; Han et al., Nature 349:697-700, 1991) by deletion of the adeno origin of replication, M13 origin of replication and sup F selection marker. An ampicillin resistance marker is then added in order to facilitate selection of the plasmid.

A full-length rat IL-1 Type 3 receptor clone in pBluescriptSK- is isolated from the phage clone described above, and cut with *EcoRV* and *HindIII*, releasing two

inserts. The inserts are then isolated and ligated to pCDM7-Amp which had been similarly cut. The resulting product is used to transform *E. coli* DH5 $\alpha$ , and colonies are examined by restriction digests for correct orientation of the two inserts (i.e., proper formation of the IL-1 Type 3R coding sequence.)

5 COS-7 (ATCC No. CRL 1651) cells are then transfected with pCDM7-Amp containing IL-1 Type 3 receptor cDNA (10  $\mu$ g DNA/10 cm plate of cells) utilizing 400  $\mu$ g/ml of DEAE-Dextran and 100  $\mu$ M chloroquine. The cells are transfected for 4 hours, then shocked with 10% DMSO for 2 minutes. The cells are then washed, and grown in DMEM containing 10% Fetal Bovine Serum for 2 days in a 24-well plate.

10

B. Expression of Human Interleukin-1 Type 3 Receptor

A full-length human IL-1 Type 3 receptor clone in pBluescriptSK- is isolated from the phage clone described above, and cut with *Not*I and *Xho*I, releasing the insert. The insert is then isolated and ligated to pCDM7-Amp which had been similarly cut. The resulting product is used to transform *E. coli* DH5 $\alpha$ , from which larger quantities of plasmid DNA may be isolated.

15 COS-7 (ATCC No. CRL 1651) cells are then transfected with pCDM7-Amp containing IL-1 Type 3 receptor cDNA (10  $\mu$ g DNA/10 cm plate of cells) utilizing 400  $\mu$ g/ml of DEAE-Dextran and 100  $\mu$ M chloroquine. The cells are transfected for 4 hours, then shocked with 10% DMSO for 2 minutes. The cells are then washed, and grown in DMEM containing 10% Fetal Bovine Serum for 2 days in a 24-well plate.

EXAMPLE 3

25 CONSTRUCTION AND EXPRESSION OF SOLUBLE  
HUMAN INTERLEUKIN-1 TYPE 3 RECEPTOR

A. Plasmid Construction

1. Vector Preparation

30 An expression vector containing the N-terminal portion of the human IL-1 type 3 receptor, also referred to as the "soluble" form of the receptor, is constructed essentially as described below. Briefly, pCDM7amp DNA (as described above) is subjected to restriction endonuclease digestion with two enzymes, *Not*I and *Xho*I, each of which have one recognition site in this vector, both located in the MCS. The product is a linearized DNA fragment with the CMV promoter/enhancer immediately upstream of the cut site, and the polyadenylation signal downstream of the cut site.

35



After digestion, the cleaved vector is isolated by agarose gel electrophoresis and purified using the Gene Clean procedure (Bio 101, San Diego, CA). The vector is now ready to combine with a DNA fragment encoding the soluble human IL-1 type 3 receptors.

5

## 2. Insert Preparation

Into this prepared vector is ligated a DNA fragment containing the coding region of the first 336 amino acids of the human IL-1 type 3 receptor set forth in Sequence ID No. 1 (from nucleotide number 129 to nucleotide number 1136).

10

Briefly, two oligonucleotides are first synthesized for use as primers in PCR. These oligonucleotides can be synthesized on a DNA synthesizer. The first primer consists of the sequence 5'-CCTACTCGAG ATGTGGTCCT TGCTGCTC-3' (Sequence ID No: 8). The first four nucleotides of this sequence serve as a spacer, and increase the efficiency of endonuclease cleavage in a subsequent reaction to be described. Nucleotides 5 through 10 encode a XhoI endonuclease cleavage site, and nucleotides 11 through 28 are identical to the N-terminal coding region of the human IL-type 3 receptor (nucleotides 129 to 146 in Sequence ID No: 1)).

The second primer consists of the sequence 5'-ATGCGCGGCC GCCTATCGAA AATCCGGAGC TGG-3' (Sequence Id No: 9). The first four nucleotides of this sequence serve as a spacer, and increase efficiency of endonuclease cleavage in a subsequent reaction to be described. Nucleotides 5 through 12 encode a *NotI* endonuclease cleavage site. Nucleotides 13 through 15 encode a translation stop codon, and nucleotides 16 through 33 are complementary to the coding region of the human IL-1 type 3 receptor immediately preceding the transmembrane region (nucleotides 1133 through 1116 in Sequence ID No. 1).

The fragment encoding soluble human IL-1 type 3 receptor is then generated by PCR. Briefly, 100ng of each primer are combined in a 0.5ml test tube, along with 1ng of the entire human IL-1 type 3 receptor DNA sequence contained in a cloning vector, such as Bluescript (Stratagene, La Jolla, CA). Ten microliters of 10X PCR buffer, 5ul of 25mM MgCl, 1ul of 25mM aTP, and 1ul of Taq polymerase/Vent polymerase (16:1 ratio) are also added to the reaction. The complete sample is then overlaid with 100ul of mineral oil to prevent evaporation, and the sample is placed in a thermocycler. Reaction conditions are: 94°C for 15 seconds, 55°C for 60 seconds, and 72°C for 60 seconds. These conditions are repeated for 25 cycles.

35

Product from the reaction is analyzed by agarose gel electrophoresis to verify the size of the fragment (1009 bp) and also to determine the approximate amount of DNA generated. The DNA is then isolated by phenol/chloroform extraction and

purified over a G-50 mini-spin column (Boehringer Mannheim, Indianapolis, IN). Approximately 10ug of the purified DNA fragment is digested with 20 units each of XhoI and NotI restriction endonucleases in a standard reaction to generate cohesive ends on the fragment which are compatible with the pCDM7 vector prepared as detailed above. The digested fragment is then agarose gel purified to remove impurities and contaminating DNA species.

### 3. Ligation

One hundred nanograms of vector DNA is combined with 100ng of insert DNA in a 1.5ml mini-tube with 1ul of 10X ligation buffer, 1ul of DNA ligase (Boehringer Mannheim), and water to a total volume of 10ul. This sample is incubated at 23°C for 2 hours.

### 4. Transformation

One hundred microliters of competent *E. coli* bacteria cells are combined with the ligation product and incubated on ice for 30 minutes. The sample is then incubated at 42°C for 45 seconds. One milliliter of bacterial medium (Circle Grow, Bio 101, San Diego, CA) is then added, and the sample is shaken at 37°C for 60 minutes. The sample is then plated on a bacterial growth plate containing bacterial medium and ampicillin at 100ug/ml (Fisher Scientific), and incubated for 16 hours at 37°C.

### 5. Construct verification

Ten colonies from the ampicillin plate are selected and grown in 1 ml of bacterial medium for 24 hours. One hundred microliters of each culture is stored by adding an equal volume of 50% glycerol solution and frozen at -70°C in mini-tubes. Plasmid DNA is then extracted from the remaining cultures by the mini-prep procedure essentially as described by Maniatis et al. (*supra*), and the recovered DNAs are analyzed by restriction digest with XhoI and NotI restriction endonucleases. The products of restriction digest are visualized by agarose gel electrophoresis and ethidium bromide staining. Correct plasmids will yield two bands: a vector band of approximately 3 kilobases, and an insert fragment of 1009 bases.

The frozen stock of a colony containing the correct plasmid is used to inoculate one liter of bacterial growth medium containing ampicillin (100ug/ml). The culture is shaken at 37°C for 24 hours, and plasmid DNA is isolated by a maxi-prep procedure (Promega). The portion on this plasmid coding for soluble human IL-1 type 3 receptor is analyzed by DNA sequencing (US Biochemical) in order to verify that the sequence is correct.

### B. Transfection Procedure and Expression

COS-7 (ATCC No. CRL 1651) or L-tk<sup>-</sup> cells (ATCC No. CCL 1.3 ) are seeded at  $1 \times 10^6$  or  $3 \times 10^6$  cells on 10 cm tissue culture dishes and incubated over night.

5 Cells are then transfected by a standard DEAE dextran method. Briefly, 10 $\mu$ g of IL-1 type 3 receptor expression plasmid DNA are diluted in 3 ml of Dulbecco's modified Minimum Essential Medium (D-MEM) supplemented with glutamine, pyruvate, 25mM HEPES, 100 microgram/ml DEAE dextran (0.5 Md., Sigma, St. Louis) and 0.1 mM chloroquine (Sigma). Cells are incubated in this transfection mixture for 4 hours at

10 37°C. After one washing step with D-MEM cells are incubated for 48 hours in D-MEM supplemented with 10% fetal calf serum. At this stage cells are ready for further analysis of the expressed IL-1 type 3 receptor.

15

### EXAMPLE 4

#### SIGNALING OF IL-1 VIA THE IL-1 TYPE 3 RECEPTOR IN A FUNCTIONAL ASSAY

IL-1 type 1 receptor cDNA and type 3 receptor cDNA are separately

20 transfected into Jurkat cells (ATCC no. TIB 152) together with a reporter plasmid consisting of the HIV promoter region (HIV-LTR) linked to the bacterial chloramphenicol acetyltransferase (CAT) gene. Stimulation of the transfected cells with human IL-1 alpha leads through a signaling cascade involving the transcription factor NF-kappaB to the production of CAT, which in turn can be measured by commercially

25 available assays (Promega, Madison, WI) (see also Leung et al., *J. Biol. Chem.* 269:1579-1582, 1994).

Results are shown in Figure 3. Briefly, approximately equal stimulation of CAT activity for both receptors can be seen over mock transfected control cells. This indicates that human IL-1 alpha can signal through the IL-1 type 3 receptor.

30

### EXAMPLE 5

#### EXPRESSION, LOCALIZATION, AND ACTIVITY OF THE IL-1 TYPE 3 RECEPTOR

35

### A. Expression Pattern of the IL-1 Type 3 Receptor

In order to determine in which rat tissues and parts of the rat brain the IL-1 Type 3 receptor is expressed, RNA protection assays are performed. Briefly, total

RNA is isolated from each tissue or part of the brain and annealed at 65°C to <sup>32</sup>P-labeled RNA generated from a plasmid containing a 600 bp fragment which covers the entire transmembrane region and portions of the extracellular and intracellular domains of the Type 3 receptor cDNA. Samples are then digested with RNase and fractionated on a denaturing polyacrylamide gel. The gel is then dried and the radioactivity quantitated using a PhosphorImager (Figure 4).

As can be seen in Figure 4, the highest level of expression is in the lung, followed by the epididymus and testis. When various areas of the brain are examined, the cerebral cortex contains the highest level of the Type 3 receptor, although other areas of the brain were also positive.

#### B. Localization of the IL-1 Type 3 Receptor by In Situ Hybridization

Utilizing *in situ* hybridization histochemistry, the IL-1 type 3 receptor may be found in the thymus and the spleen. In the thymus the signal is most prominent in the cortical region and not in the medulla. Within the rat brain the IL-1 type 3 receptor expression is detectable in the hippocampus and the fourth ventricle. This is in contrast to the localization of the IL-1 type 1 receptor which is restricted to the dentate gyrus granule cells.

Briefly, dissected tissue is frozen in isopentane cooled to -42°C and subsequently stored at -80°C prior to sectioning on a cryostat. Slide-mounted tissue sections are then stored at -80°C. Sections are removed from storage and placed directly into 4% buffered paraformaldehyde at room temperature. After 60 minutes, slides are rinsed in isotonic phosphate buffered saline (10 min.) and treated with proteinase K (1 µg/ml in 100 mM Tris/HCl, pH 8.0) for 10 minutes at 37°C. Subsequently, sections are successively washed in water (1 min.), 0.1 M triethanolamine (pH 8.0, plus 0.25% acetic anhydride) for 10 minutes and 2X SSC (0.3 mM NaCl, 0.03 mM sodium citrate, pH 7.2) for 5 minutes. Sections are then dehydrated through graded alcohols and air dried. Post-fixed sections are hybridized with 1.0 x 10<sup>6</sup> dpm [35S]UTP-labeled riboprobes in hybridization buffer containing 75% formamide, 10% dextran sulphate, 3X SSC, 50 mM sodium phosphate buffer pH 7.4, 1X Denhardt's solution, 0.1 mg/ml yeast tRNA and 10 mM dithiothreitol in a total volume of 30 µl. The diluted probe is applied to sections on a glass coverslip and hybridized overnight at 55°C in a humid environment. Post-hybridization, sections are washed in 2X SSC for 5 minutes and then treated with RNase A (200 µg/ml in 10 mM Tris/HCl, pH 8.0, containing 0.5 M NaCl) for 60 minutes at 37°C. Subsequently, sections are washed in 2X SSC for 5 minutes, 1X SSC for 5 minutes, 0.1X SSC for 60 minutes at 70°C, 0.5X

SSC at room temperature for 5 minutes and then dehydrated in graded alcohols and air dried. For signal detection, sections are placed on Kodak Bio Max X-ray film and exposed for the required length of time or dipped in photographic emulsion (Amersham LM-1) for high resolution analysis. Autoradiograms are analyzed using automated  
5 image analysis (DAGE camera/Mac II) while dipped sections were examined using a Zeiss Axioscope.

C. Inhibition of Thymocyte Proliferation by the IL-1 Type 3 Receptor

10 Ability of the IL-1 type 3 receptor to inhibit mouse thymocyte proliferation may also be examined. Briefly, the proliferative response of T lymphocyte lectins such as phytohemagglutinin (PHA) is very low, but is markedly enhanced by IL-1. Thus, soluble type human and rat type 3 receptors may be utilized to competitively inhibit proliferation of mouse thymocytes stimulated by IL-1. Soluble human Type 1  
15 receptor produced in baculovirus may be used as a positive control.

Briefly, soluble IL-1 type 1 or type 3 receptors are added to wells of a 96 well plate and serially diluted IL-1 is also added. Thymi are removed from young mice and a single cell suspension prepared in tissue culture media. Cells are washed 3 times and resuspended at a concentration of  $10^7$  cells/ml. Cells are plated at 100 microliters in  
20 a 96 well flat bottom microtiter plate. PHA is added to stimulate the cells. Plates are then incubated for 48 hours in a 37°C, 5% CO<sub>2</sub> humidified incubator, and [<sup>3</sup>H] thymidine is added to the cells for the last 4 to 6 hours. Cells are then harvested and the [<sup>3</sup>H] thymidine incorporation determined by liquid scintillation counting.

As shown in Figure 5, both human IL-1 type 3 and rat IL-1 type 3  
25 receptors effectively inhibit thymocyte proliferation in a manner similar to that observed for soluble human type 1 receptor. This result strongly indicates that the type 3 receptor inhibits thymocyte proliferation by binding to the exogenously added IL-1.

From the foregoing, it will be appreciated that, although specific  
30 embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(i) APPLICANTS: Lovenberg, Timothy W.  
Oltersdorf, Tilman  
Liaw, Chen W.  
Clevenger, William  
DeSouza, Errol B.

(ii) TITLE OF INVENTION: INTERLEUKIN-1 TYPE 3 RECEPTORS

(iii) NUMBER OF SEQUENCES: 9

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Seed and Berry  
(B) STREET: 6300 Columbia Center, 701 Fifth Avenue  
(C) CITY: Seattle  
(D) STATE: Washington  
(E) COUNTRY: US  
(F) ZIP: 98104

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: McMasters, David D.  
(B) REGISTRATION NUMBER: 33.963

(C) REFERENCE/DOCKET NUMBER: 690068.402PC

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (206) 622-4900

(B) TELEFAX: (206) 682-6031

(C) TELEX: 3723836

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1965 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 129..1814

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGCCCGCCCA CGGCGGCGGG GAAATACCTA GGCATGGAAG TGGCATGACA GGGCTCGTGT	60
CCCTGTCATA TTTTCCACTC TCCACGAGGT CCTGCGCGCT TCAATCCTGC AGGCAGCCCCG	120
GTTTGGGG ATG TGG TCC TTG CTG CTC TGC GGG TTG TCC ATC GCC CTT CCA	170
Met Trp Ser Leu Leu Leu Cys Gly Leu Ser Ile Ala Leu Pro	
1 5 10	
CTG TCT GTC ACA GCA GAT GGA TGC AAG GAC ATT TTT ATG AAA AAT GAG	218
Leu Ser Val Thr Ala Asp Gly Cys Lys Asp Ile Phe Met Lys Asn Glu	
15 20 25 30	
ATA CTT TCA GCA AGC CAG CCT TTT GCT TTT AAT TGT ACA TTC CCT CCC	266

ATC CCA GTG TCC AAA ATC ATA CAG TCT AGA ATT CAC CAG GAC GAG ACT 362  
Ile Pro Val Ser Lys Ile Ile Gln Ser Arg Ile His Gln Asp Glu Thr  
65 70 75

TGT GTT ATA AAG GGT AGA GAC AGC TGT CAT AGA ATA CAT GTA AAC CTA 458  
Cys Val Ile Lys Gly Arg Asp Ser Cys His Arg Ile His Val Asn Leu  
95 100 105 110

AAT TTA TCA GAT GAG TAC AAG CAA ATA TTA CAT CTT GGA AAA GAT GAT  
Asn Leu Ser Asp Glu Tyr Lys Gln Ile Leu His Leu Gly Lys Asp Asp  
130 135 140

ATA AAG TGG TAT AAG GAC TGT AAC GAG ATT AAA GGG GAG CGG TTC ACT 650  
Ile Lys Trp Tyr Lys Asp Cys Asn Glu Ile Lys Gly Glu Arg Phe Thr  
160 165 170

GTT TTG GAA ACC AGG CTT TTG GTG AGC AAT GTC TCG GCA GAG GAC AGA  
Val Leu Glu Thr Arg Leu Leu Val Ser Asn Val Ser Ala Glu Asp Arg

175	180	185	190	
GGG AAC TAC GCG TGT CAA GCC ATA CTG ACA CAC TCA GGG AAG CAG TAC				746
Gly Asn Tyr Ala Cys Gln Ala Ile Leu Thr His Ser Gly Lys Gln Tyr				
195	200	205		
GAG GTT TTA AAT GGC ATC ACT GTG AGC ATT ACA GAA AGA GCT GGA TAT				794
Glu Val Leu Asn Gly Ile Thr Val Ser Ile Thr Glu Arg Ala Gly Tyr				
210	215	220		
GGA GGA AGT GTC CCT AAA ATC ATT TAT CCA AAA AAT CAT TCA ATT GAA				842
Gly Gly Ser Val Pro Lys Ile Ile Tyr Pro Lys Asn His Ser Ile Glu				
225	230	235		
GTA CAG CTT GGT ACC ACT CTG ATT GTG GAC TGC AAT GTA ACA GAC ACC				890
Val Gln Leu Gly Thr Thr Leu Ile Val Asp Cys Asn Val Thr Asp Thr				
240	245	250		
AAG GAT AAT ACA AAT CTA CGA TGC TGG AGA GTC AAT AAC ACT TTG GTG				938
Lys Asp Asn Thr Asn Leu Arg Cys Trp Arg Val Asn Asn Thr Leu Val				
255	260	265	270	
GAT GAT TAC TAT GAT GAA TCC AAA CGA ATC AGA GAA GGG GTG GAA ACC				986
Asp Asp Tyr Tyr Asp Glu Ser Lys Arg Ile Arg Glu Gly Val Glu Thr				
275	280	285		
CAT GTC TCT TTT CGG GAA CAT AAT TTG TAC ACA GTA AAC ATC ACC TTC				1034
His Val Ser Phe Arg Glu His Asn Leu Tyr Thr Val Asn Ile Thr Phe				
290	295	300		
TTG GAA GTG AAA ATG GAA GAT TAT GGC CTT CCT TTC ATG TGC CAC GCT				1082
Leu Glu Val Lys Met Glu Asp Tyr Gly Leu Pro Phe Met Cys His Ala				
305	310	315		
GGA GTG TCC ACA GCA TAC ATT ATA TTA CAG CTC CCA GCT CCG GAT TTT				1130
Gly Val Ser Thr Ala Tyr Ile Ile Leu Gln Leu Pro Ala Pro Asp Phe				
320	325	330		

CGA GCT TAC TTG ATA GGA GGG CTT ATC GCC TTG GTG GCT GTG GCT GTG	1178
Arg Ala Tyr Leu Ile Gly Gly Leu Ile Ala Leu Val Ala Val Ala Val	
335 340 345 350	
TCT GTT GTG TAC ATA TAC AAC ATT TTT AAG ATC GAC ATT GTT CTT TGG	1226
Ser Val Val Tyr Ile Tyr Asn Ile Phe Lys Ile Asp Ile Val Leu Trp	
355 360 365	
TAT CGA AGT GCC TTC CAT TCT ACA GAG ACC ATA GTA GAT GGG AAG CTG	1274
Tyr Arg Ser Ala Phe His Ser Thr Glu Thr Ile Val Asp Gly Lys Leu	
370 375 380	
TAT GAC GCC TAT GTC TTA TAC CCC AAG CCC CAC AAG GAA AGC CAG AGG	1322
Tyr Asp Ala Tyr Val Leu Tyr Pro Lys Pro His Lys Glu Ser Gln Arg	
385 390 395	
CAT GCC GTG GAT GCC CTG GTG TTG AAT ATC CTG CCC GAG GTG TTG GAG	1370
His Ala Val Asp Ala Leu Val Leu Asn Ile Leu Pro Glu Val Leu Glu	
400 405 410	
AGA CAA TGT GGA TAT AAG TTG TTT ATA TTC GGC AGA GAT GAA TTC CCT	1418
Arg Gln Cys Gly Tyr Lys Leu Phe Ile Phe Gly Arg Asp Glu Phe Pro	
415 420 425 430	
GGA CAA GCC GTG GCC AAT GTC ATC GAT GAA AAC GTT AAG CTG TGC AGG	1466
Gly Gln Ala Val Ala Asn Val Ile Asp Glu Asn Val Lys Leu Cys Arg	
435 440 445	
AGG CTG ATT GTC ATT GTG GTC CCC GAA TCG CTG GGC TTT GGC CTG TTG	1514
Arg Leu Ile Val Ile Val Val Pro Glu Ser Leu Gly Phe Gly Leu Leu	
450 455 460	
AAG AAC CTG TCA GAA GAA CAA ATC GCG GTC TAC AGT GCC CTG ATC CAG	1562
Lys Asn Leu Ser Glu Glu Gln Ile Ala Val Tyr Ser Ala Leu Ile Gln	
465 470 475	



GAC GGG ATG AAG GTT ATT CTC ATT GAG CTG GAG AAA ATC GAG GAC TAC 1610  
Asp Gly Met Lys Val Ile Leu Ile Glu Leu Glu Lys Ile Glu Asp Tyr  
480 485 490

ACA GTC ATG CCA GAG TCA ATT CAG TAC ATC AAA CAG AAG CAT GGT GCC 1658  
Thr Val Met Pro Glu Ser Ile Gln Tyr Ile Lys Gln Lys His Gly Ala  
495 500 505 510

ATC CGG TGG CAT GGG GAC TTC ACG GAG CAG TCA CAG TGT ATG AAG ACC 1706  
Ile Arg Trp His Gly Asp Phe Thr Glu Gln Ser Gln Cys Met Lys Thr  
515 520 525

AAG TTT TGG AAG ACA GTG AGA TAC CAC ATG CCG CCC AGA AGG TGT CGG 1754  
Lys Phe Trp Lys Thr Val Arg Tyr His Met Pro Pro Arg Arg Cys Arg  
530 535 540

CCG TTT CTC CGG TCC ACG TGC CGC AGC ACA CAC CTC TGT ACC GCA CCG 1802  
Pro Phe Leu Arg Ser Thr Cys Arg Ser Thr His Leu Cys Thr Ala Pro  
545 550 555

CAG GCC CAG AAC TAGGCTCAAG AAGAAAGAAG TGTACTCTCA CGACTGGCTA 1854  
Gln Ala Gln Asn  
560

AGACTTGCTG GACTGACACC TATGGCTGGA AGATGACTTG TTTTGCTCCA TGTCTCCTCA 1914

TTCCTACACC TATTTTCTGC TGCAGGATGA GGCTAGGGTT AGCATTCTAG A 1965

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 562 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Trp Ser Leu Leu Leu Cys Gly Leu Ser Ile Ala Leu Pro Leu Ser  
1 5 10 15

Val Thr Ala Asp Gly Cys Lys Asp Ile Phe Met Lys Asn Glu Ile Leu  
20 25 30

Ser Ala Ser Gln Pro Phe Ala Phe Asn Cys Thr Phe Pro Pro Ile Thr  
35 40 45

Ser Gly Glu Val Ser Val Thr Trp Tyr Lys Asn Ser Ser Lys Ile Pro  
50 55 60

Val Ser Lys Ile Ile Gln Ser Arg Ile His Gln Asp Glu Thr Trp Ile  
65 70 75 80

Leu Phe Leu Pro Met Glu Trp Gly Asp Ser Gly Val Tyr Gln Cys Val  
85 90 95

Ile Lys Gly Arg Asp Ser Cys His Arg Ile His Val Asn Leu Thr Val  
100 105 110

Phe Glu Lys His Trp Cys Asp Thr Ser Ile Gly Gly Leu Pro Asn Leu  
115 120 125

Ser Asp Glu Tyr Lys Gln Ile Leu His Leu Gly Lys Asp Asp Ser Leu  
130 135 140

Thr Cys His Leu His Phe Pro Lys Ser Cys Val Leu Gly Pro Ile Lys  
145 150 155 160

Trp Tyr Lys Asp Cys Asn Glu Ile Lys Gly Glu Arg Phe Thr Val Leu  
165 170 175

Glu Thr Arg Leu Leu Val Ser Asn Val Ser Ala Glu Asp Arg Gly Asn

180 185 190  
Tyr Ala Cys Gln Ala Ile Leu Thr His Ser Gly Lys Gln Tyr Glu Val  
195 200 205  
Leu Asn Gly Ile Thr Val Ser Ile Thr Glu Arg Ala Gly Tyr Gly Gly  
210 215 220  
Ser Val Pro Lys Ile Ile Tyr Pro Lys Asn His Ser Ile Glu Val Gln  
225 230 235 240  
Leu Gly Thr Thr Leu Ile Val Asp Cys Asn Val Thr Asp Thr Lys Asp  
245 250 255  
Asn Thr Asn Leu Arg Cys Trp Arg Val Asn Asn Thr Leu Val Asp Asp  
260 265 270  
Tyr Tyr Asp Glu Ser Lys Arg Ile Arg Glu Gly Val Glu Thr His Val  
275 280 285  
Ser Phe Arg Glu His Asn Leu Tyr Thr Val Asn Ile Thr Phe Leu Glu  
290 295 300  
Val Lys Met Glu Asp Tyr Gly Leu Pro Phe Met Cys His Ala Gly Val  
305 310 315 320  
Ser Thr Ala Tyr Ile Ile Leu Gln Leu Pro Ala Pro Asp Phe Arg Ala  
325 330 335  
Tyr Leu Ile Gly Gly Leu Ile Ala Leu Val Ala Val Ala Val Ser Val  
340 345 350  
Val Tyr Ile Tyr Asn Ile Phe Lys Ile Asp Ile Val Leu Trp Tyr Arg  
355 360 365  
Ser Ala Phe His Ser Thr Glu Thr Ile Val Asp Gly Lys Leu Tyr Asp  
370 375 380

Ala Tyr Val Leu Tyr Pro Lys Pro His Lys Glu Ser Gln Arg His Ala  
 385 390 395 400

Val Asp Ala Leu Val Leu Asn Ile Leu Pro Glu Val Leu Glu Arg Gln  
 405 410 415

Cys Gly Tyr Lys Leu Phe Ile Phe Gly Arg Asp Glu Phe Pro Gly Gln  
 420 425 430

Ala Val Ala Asn Val Ile Asp Glu Asn Val Lys Leu Cys Arg Arg Leu  
 435 440 445

Ile Val Ile Val Val Pro Glu Ser Leu Gly Phe Gly Leu Leu Lys Asn  
 450 455 460

Leu Ser Glu Glu Gln Ile Ala Val Tyr Ser Ala Leu Ile Gln Asp Gly  
 465 470 475 480

Met Lys Val Ile Leu Ile Glu Leu Glu Lys Ile Glu Asp Tyr Thr Val  
 485 490 495

Met Pro Glu Ser Ile Gln Tyr Ile Lys Gln Lys His Gly Ala Ile Arg  
 500 505 510

Trp His Gly Asp Phe Thr Glu Gln Ser Gln Cys Met Lys Thr Lys Phe  
 515 520 525

Trp Lys Thr Val Arg Tyr His Met Pro Pro Arg Arg Cys Arg Pro Phe  
 530 535 540

Leu Arg Ser Thr Cys Arg Ser Thr His Leu Cys Thr Ala Pro Gln Ala  
 545 550 555 560

Gln Asn

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2044 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 89..1771

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

CCGGCTGGCC TAGGATCAGG CAAGAAAAGG CTGAACGCCT TTCTAAGGAC GGACTCTTTC      60
TGTACAGCTC CACTTGGGGA AGCCCGAA ATG GGG ATG CCA CCC TTG CTC TTC      112
      Met Gly Met Pro Pro Leu Leu Phe
              1              5
TGT TGG GTG TCT TTC GTG CTT CCA CTT TTT GTG GCA GCA GGT AAC TGT      160
Cys Trp Val Ser Phe Val Leu Pro Leu Phe Val Ala Ala Gly Asn Cys
      10              15              20
ACT GAT GTC TAT ATG CAC CAT GAG ATG ATT TCA GAG GGC CAG CCT TTC      208
Thr Asp Val Tyr Met His His Glu Met Ile Ser Glu Gly Gln Pro Phe
      25              30              35              40
CCC TTC AAC TGC ACA TAC CCT CCA GTA ACA AAC GGG GCA GTG AAT CTG      256
Pro Phe Asn Cys Thr Tyr Pro Pro Val Thr Asn Gly Ala Val Asn Leu
              45              50              55
ACA TGG CAT AGA ACA CCC AGT AAG AGC CCA ATC TCC ATC AAC AGA CAC      304
Thr Trp His Arg Thr Pro Ser Lys Ser Pro Ile Ser Ile Asn Arg His
      60              65              70

```



GTT AGA ATT CAC CAG GAC CAG TCC TGG ATT TTG TTT CTT CCG TTG GCA Val Arg Ile His Gln Asp Gln Ser Trp Ile Leu Phe Leu Pro Leu Ala 75 80 85	352
TTG GAG GAC TCA GGC ATC TAT CAA TGT GTT ATA AAG GAT GCC CAC AGC Leu Glu Asp Ser Gly Ile Tyr Gln Cys Val Ile Lys Asp Ala His Ser 90 95 100	400
TGT TAC CGA ATA GCT ATA AAC CTA ACC GTT TTT AGA AAA CAC TGG TGC Cys Tyr Arg Ile Ala Ile Asn Leu Thr Val Phe Arg Lys His Trp Cys 105 110 115 120	448
GAC TCT TCC AAC GAA GAG AGT TCC ATA AAT TCC TCA GAT GAG TAC CAG Asp Ser Ser Asn Glu Glu Ser Ser Ile Asn Ser Ser Asp Glu Tyr Gln 125 130 135	496
CAA TGG TTA CCC ATA GGA AAA TCG GGC AGT CTG ACG TGC CAT CTC TAC Gln Trp Leu Pro Ile Gly Lys Ser Gly Ser Leu Thr Cys His Leu Tyr 140 145 150	544
TTC CCA GAG AGC TGT GTT TTG GAT TCA ATA AAG TGG TAT AAG GGT TGT Phe Pro Glu Ser Cys Val Leu Asp Ser Ile Lys Trp Tyr Lys Gly Cys 155 160 165	592
GAA GAG ATT AAA GTG AGC AAG AAG TTT TGC CCT ACA GGA ACA AAG CTT Glu Glu Ile Lys Val Ser Lys Lys Phe Cys Pro Thr Gly Thr Lys Leu 170 175 180	640
CTT GTT AAC AAC ATC GAC GTG GAG GAT AGT GGG AGC TAT GCA TGC TCA Leu Val Asn Asn Ile Asp Val Glu Asp Ser Gly Ser Tyr Ala Cys Ser 185 190 195 200	688
GCC AGA CTG ACA CAC TTG GGG AGA ATC TTC ACG GTT AGA AAC TAC ATT Ala Arg Leu Thr His Leu Gly Arg Ile Phe Thr Val Arg Asn Tyr Ile 205 210 215	736

GCT GTG AAT ACC AAG GAA GTT GGG TCT GGA GGA AGG ATC CCT AAC ATC 784  
 Ala Val Asn Thr Lys Glu Val Gly Ser Gly Gly Arg Ile Pro Asn Ile  
                   220                  225                  230

ACG TAT CCA AAA AAC AAC TCC ATT GAA GTT CAA CTT GGC TCC ACC CTC 832  
 Thr Tyr Pro Lys Asn Asn Ser Ile Glu Val Gln Leu Gly Ser Thr Leu  
                   235                  240                  245

ATT GTG GAC TGC AAT ATA ACA GAC ACG AAG GAG AAT ACG AAC CTC AGA 880  
 Ile Val Asp Cys Asn Ile Thr Asp Thr Lys Glu Asn Thr Asn Leu Arg  
                   250                  255                  260

TGC TGG CGA GTT AAC AAC ACC CTG GTG GAC GAT TAC TAC AAC GAC TTC 928  
 Cys Trp Arg Val Asn Asn Thr Leu Val Asp Asp Tyr Tyr Asn Asp Phe  
                   265                  270                  275                  280

AAA CGC ATC CAG GAA GGA ATC GAA ACC AAT CTG TCT CTG AGG AAT CAC 976  
 Lys Arg Ile Gln Glu Gly Ile Glu Thr Asn Leu Ser Leu Arg Asn His  
                   285                  290                  295

ATT CTG TAC ACA GTG AAC ATA ACA TTC TTA GAA GTG AAA ATG GAG GAC 1024  
 Ile Leu Tyr Thr Val Asn Ile Thr Phe Leu Glu Val Lys Met Glu Asp  
                   300                  305                  310

TAC GGC CAT CCT TTC ACA TGC CAC GCT GCG GTG TCC GCA GCC TAC ATC 1072  
 Tyr Gly His Pro Phe Thr Cys His Ala Ala Val Ser Ala Ala Tyr Ile  
                   315                  320                  325

ATT CTG AAA CGC CCA GCT CCA GAC TTC CGG GCT TAC CTC ATA GGA GGT 1120  
 Ile Leu Lys Arg Pro Ala Pro Asp Phe Arg Ala Tyr Leu Ile Gly Gly  
                   330                  335                  340

CTC ATG GCT TTC CTA CTT CTG GCC GTG TCC ATT CTG TAC ATC TAC AAC 1168  
 Leu Met Ala Phe Leu Leu Leu Ala Val Ser Ile Leu Tyr Ile Tyr Asn  
                   345                  350                  355                  360

ACC TTT AAG GTC GAC ATC GTG CTT TGG TAT AGG AGT ACC TTC CAC ACT 1216

Thr Phe Lys Val Asp Ile Val Leu Trp Tyr Arg Ser Thr Phe His Thr	365	370	375	
GCC CAG GCT CCA GAT GAC GAG AAG CTG TAT GAT GCC TAT GTC TTA TAC				1264
Ala Gln Ala Pro Asp Asp Glu Lys Leu Tyr Asp Ala Tyr Val Leu Tyr	380	385	390	
CCC AAG TAC CCA AGA GAA AGC CAG GGC CAT GAT GTG GAC ACA CTG GTG				1312
Pro Lys Tyr Pro Arg Glu Ser Gln Gly His Asp Val Asp Thr Leu Val	395	400	405	
TTG AAG ATC TTG CCC GAG GTG CTG GAG AAA CAG TGT GGA TAT AAG TTA				1360
Leu Lys Ile Leu Pro Glu Val Leu Glu Lys Gln Cys Gly Tyr Lys Leu	410	415	420	
TTC ATA TTT GGC AGG GAT GAA TTC CCT GGA CAA GCT GTG GCC AGC GTC				1408
Phe Ile Phe Gly Arg Asp Glu Phe Pro Gly Gln Ala Val Ala Ser Val	425	430	435	440
ATT GAT GAA AAC ATT AAG CTG TGT AGG AGG CTG ATG GTC CTC GTG GCA				1456
Ile Asp Glu Asn Ile Lys Leu Cys Arg Arg Leu Met Val Leu Val Ala	445	450	455	
CCA GAG ACA TCC AGC TTC AGC TTT CTG AAG AAC TTG ACT GAA GAA CAA				1504
Pro Glu Thr Ser Ser Phe Ser Phe Leu Lys Asn Leu Thr Glu Glu Gln	460	465	470	
ATC GCT GTC TAC AAT GCC CTC GTC CAG GAC GGC ATG AAG GTC ATT CTG				1552
Ile Ala Val Tyr Asn Ala Leu Val Gln Asp Gly Met Lys Val Ile Leu	475	480	485	
ATT GAA CTG GAG AGA GTC AAG GAC TAC AGC ACC ATG CCC GAG TCC ATT				1600
Ile Glu Leu Glu Arg Val Lys Asp Tyr Ser Thr Met Pro Glu Ser Ile	490	495	500	
CAG TAC ATC CGA CAG AAG CAC GGG GCC ATC CAG TGG GAT GGG GAC TTC				1648
Gln Tyr Ile Arg Gln Lys His Gly Ala Ile Gln Trp Asp Gly Asp Phe				

505	510	515	520	
ACA GAG CAG GCA CAG TGC GCC AAG ACG AAA TTC TGG AAG AAA GTG AGA				1696
Thr Glu Gln Ala Gln Cys Ala Lys Thr Lys Phe Trp Lys Lys Val Arg				
	525	530	535	
TAT CAT ATG CCA CCC AGG AGG TAC CCG GCA TCT CCC CCC GTC CAG CTG				1744
Tyr His Met Pro Pro Arg Arg Tyr Pro Ala Ser Pro Pro Val Gln Leu				
	540	545	550	
CTA GGA CAC ACA CCC CGC ATA CCA GGC TAGTGCAGTG CCACCGCCAC				1791
Leu Gly His Thr Pro Arg Ile Pro Gly				
	555	560		
GGGGCTCATA ACTCCTTAAG AGCGGTTAGT GTGTGGTGGC TCGCACTACA ACCTCTCTGG				1851
ATCATCTACC CCCGTAGCTT GCTCTTTTGT GCTTGTGAGC GACCTCGTCC TTAGCCACGT				1911
CATATTTTGA TTTTGTGTTT GTTTTGTGTTG TTTGTTGTAT GCTTTTAGTC ATAGCTGATT				1971
CGTACTACTC CTGTTTGCTT CATGGTTCCT GAATCCCAGA GACTCCCTGA GCATGGGTGG				2031
CTATCATGTT GGG				2044

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 561 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Gly Met Pro Pro Leu Leu Phe Cys Trp Val Ser Phe Val Leu Pro

WO 96/07739

1 5 10 15  
 Leu Phe Val Ala Ala Gly Asn Cys Thr Asp Val Tyr Met His His Glu  
 20 25 30  
 Met Ile Ser Glu Gly Gln Pro Phe Pro Phe Asn Cys Thr Tyr Pro Pro  
 35 40 45  
 Val Thr Asn Gly Ala Val Asn Leu Thr Trp His Arg Thr Pro Ser Lys  
 50 55 60  
 Ser Pro Ile Ser Ile Asn Arg His Val Arg Ile His Gln Asp Gln Ser  
 65 70 75 80  
 Trp Ile Leu Phe Leu Pro Leu Ala Leu Glu Asp Ser Gly Ile Tyr Gln  
 85 90 95  
 Cys Val Ile Lys Asp Ala His Ser Cys Tyr Arg Ile Ala Ile Asn Leu  
 100 105 110  
 Thr Val Phe Arg Lys His Trp Cys Asp Ser Ser Asn Glu Glu Ser Ser  
 115 120 125  
 Ile Asn Ser Ser Asp Glu Tyr Gln Gln Trp Leu Pro Ile Gly Lys Ser  
 130 135 140  
 Gly Ser Leu Thr Cys His Leu Tyr Phe Pro Glu Ser Cys Val Leu Asp  
 145 150 155 160  
 Ser Ile Lys Trp Tyr Lys Gly Cys Glu Glu Ile Lys Val Ser Lys Lys  
 165 170 175  
 Phe Cys Pro Thr Gly Thr Lys Leu Leu Val Asn Asn Ile Asp Val Glu  
 180 185 190  
 Asp Ser Gly Ser Tyr Ala Cys Ser Ala Arg Leu Thr His Leu Gly Arg  
 195 200 205



Ile Phe Thr Val Arg Asn Tyr Ile Ala Val Asn Thr Lys Glu Val Gly  
210 215 220

Ser Gly Gly Arg Ile Pro Asn Ile Thr Tyr Pro Lys Asn Asn Ser Ile  
225 230 235 240

Glu Val Gln Leu Gly Ser Thr Leu Ile Val Asp Cys Asn Ile Thr Asp  
245 250 255

Thr Lys Glu Asn Thr Asn Leu Arg Cys Trp Arg Val Asn Asn Thr Leu  
260 265 270

Val Asp Asp Tyr Tyr Asn Asp Phe Lys Arg Ile Gln Glu Gly Ile Glu  
275 280 285

Thr Asn Leu Ser Leu Arg Asn His Ile Leu Tyr Thr Val Asn Ile Thr  
290 295 300

Phe Leu Glu Val Lys Met Glu Asp Tyr Gly His Pro Phe Thr Cys His  
305 310 315 320

Ala Ala Val Ser Ala Ala Tyr Ile Ile Leu Lys Arg Pro Ala Pro Asp  
325 330 335

Phe Arg Ala Tyr Leu Ile Gly Gly Leu Met Ala Phe Leu Leu Leu Ala  
340 345 350

Val Ser Ile Leu Tyr Ile Tyr Asn Thr Phe Lys Val Asp Ile Val Leu  
355 360 365

Trp Tyr Arg Ser Thr Phe His Thr Ala Gln Ala Pro Asp Asp Glu Lys  
370 375 380

Leu Tyr Asp Ala Tyr Val Leu Tyr Pro Lys Tyr Pro Arg Glu Ser Gln  
385 390 395 400

WO 96/07739

52

Gly His Asp Val Asp Thr Leu Val Leu Lys Ile Leu Pro Glu Val Leu  
 405 410 415

Glu Lys Gln Cys Gly Tyr Lys Leu Phe Ile Phe Gly Arg Asp Glu Phe  
 420 425 430

Pro Gly Gln Ala Val Ala Ser Val Ile Asp Glu Asn Ile Lys Leu Cys  
 435 440 445

Arg Arg Leu Met Val Leu Val Ala Pro Glu Thr Ser Ser Phe Ser Phe  
 450 455 460

Leu Lys Asn Leu Thr Glu Glu Gln Ile Ala Val Tyr Asn Ala Leu Val  
 465 470 475 480

Gln Asp Gly Met Lys Val Ile Leu Ile Glu Leu Glu Arg Val Lys Asp  
 485 490 495

Tyr Ser Thr Met Pro Glu Ser Ile Gln Tyr Ile Arg Gln Lys His Gly  
 500 505 510

Ala Ile Gln Trp Asp Gly Asp Phe Thr Glu Gln Ala Gln Cys Ala Lys  
 515 520 525

Thr Lys Phe Trp Lys Lys Val Arg Tyr His Met Pro Pro Arg Arg Tyr  
 530 535 540

Pro Ala Ser Pro Pro Val Gln Leu Leu Gly His Thr Pro Arg Ile Pro  
 545 550 555 560

Gly

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 8 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Asp Tyr Lys Asp Asp Asp Asp Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CTTCAACTGC ACATACCCTC CAGTAACAAA CGGGGCAGTG AATCTGACAT

50

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCTCCCATAA CATCTGGGGA AGTCAGTGTA ACATGGTATA AAAATTCTAG C

51

## (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CCTACTCGAG ATGTGGTCCT TGCTGCTC

28

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATGCGCGGCC GCCTATCGAA AATCCGGAGC TGG

33

Claims

1. An isolated nucleic acid molecule encoding an Interleukin-1 Type 3 receptor.
2. The isolated nucleic acid molecule according to claim 1, comprising the sequence of nucleotides in Sequence I.D. No. 1, from nucleotide number 129 to nucleotide number 1814.
3. The isolated nucleic acid molecule according to claim 1 wherein said molecule encodes a protein having the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 562.
4. The isolated nucleic acid molecule according to claim 1, comprising the sequence of nucleotides in Sequence I.D. No. 3, from nucleotide number 89 to nucleotide number 1771.
5. The isolated nucleic acid molecule according to claim 1 wherein said molecule encodes a protein having the amino acid sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 561.
6. The isolated nucleic acid molecule according to claim 1 wherein said molecule encodes a human Interleukin-1 Type 3 receptor.
7. The isolated nucleic acid molecule according to claim 1 wherein said molecule encodes a rat Interleukin-1 Type 3 receptor.
8. An isolated nucleic acid molecule encoding soluble Interleukin-1 Type 3 receptor.
9. The isolated nucleic acid molecule according to claim 8, comprising the sequence of nucleotides in Sequence I.D. No. 1, from nucleotide number 129 to nucleotide number 1136.

10. The isolated nucleic acid molecule according to claim 8 wherein said molecule encodes a protein having the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 336.

11. The isolated nucleic acid molecule according to claim 8, comprising the sequence of nucleotides in Sequence I.D. No. 3, from nucleotide number 89 to nucleotide number 1102.

12. The isolated nucleic acid molecule according to claim 8 wherein said molecule encodes a protein having the amino acid sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 338.

13. The isolated nucleic acid molecule according to claim 8 wherein said molecule encodes a soluble human Interleukin-1 Type 3 receptor.

14. The isolated nucleic acid molecule according to claim 8 wherein said molecule encodes a soluble rat Interleukin-1 Type 3 receptor.

15. A recombinant expression vector, comprising a promoter operably linked to a nucleic acid molecule according to any one of claims 1-14.

16. A recombinant viral vector capable of directing the expression of a nucleic acid molecule according to any one of claims 1-14 wherein said vector is selected from the group consisting of retroviral vectors, adenoviral vectors, and herpes simplex virus vectors.

17. A host cell containing a recombinant vector according to any one of claims 15 or 16.

18. An isolated Interleukin-1 Type 3 receptor.

19. The isolated Interleukin-1 Type 3 receptor according to claim 18 having the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 562.



20. The isolated Interleukin-1 Type 3 receptor according to claim 18 having the amino acid sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 561.

21. The isolated Interleukin-1 Type 3 receptor according to claim 18 wherein said receptor is a human Interleukin-1 Type 3 receptor.

22. The isolated Interleukin-1 Type 3 receptor according to claim 18 wherein said receptor is a rat Interleukin-1 Type 3 receptor.

23. An isolated soluble Interleukin-1 Type 3 receptor.

24. The isolated soluble Interleukin-1 Type 3 receptor according to claim 23 having the amino acid sequence of Sequence I.D. No. 2, from amino acid number 1 to amino acid number 336.

25. The isolated soluble Interleukin-1 Type 3 receptor according to claim 23 having the amino acid sequence of Sequence I.D. No. 4, from amino acid number 1 to amino acid number 338.

26. The isolated soluble Interleukin-1 Type 3 receptor according to claim 23 wherein said receptor is a human Interleukin-1 Type 3 receptor.

27. The isolated soluble Interleukin-1 Type 3 receptor according to claim 23 wherein said receptor is a rat Interleukin-1 Type 3 receptor.

28. An isolated antibody capable of specifically binding to an Interleukin-1 Type 3 receptor.

29. The antibody according to claim 28 wherein said antibody is selected from the group consisting of polyclonal antibodies, monoclonal antibodies, and antibody fragments.


30. The antibody according to claim 28 wherein said antibody is capable of blocking the binding of IL-1 to an Interleukin-1 Type 3 receptor.

31. The antibody according to claim 28 wherein said antibody is selected from the group consisting of murine and human antibodies.

32. A hybridoma which produces an antibody according to any one of claims 28-31.

33. A nucleic acid probe of at least 18 nucleotides in length which is capable of specifically hybridizing to a nucleic acid sequence encoding an Interleukin-1 Type 3 receptor.

1/4




*Fig. 1*

2/4

HOMOLOGY OF THE HUMAN IL-1 TYPE 3 RECEPTOR WITH  
RELATED RECEPTORS

		OVERALL	EXTRACELLULAR	INTRACELLULAR	MEMBRANE
RAT	IL-1 R3	66	63	70	66
HUMAN	IL-1 R1	42	34	55	60
RAT	IL-1 R1	40	34	52	60
MOUSE	IL-1 R1	40			
HUMAN	IL-1 R2	23			
MOUSE	IL-1 R2	23			
MOUSE	ST-2L	29			

*Fig. 2**Fig. 3*

3/4




Fig. 4

4/4




Fig. 5A




Fig. 5B

SUBSTITUTE SHEET (RULE 26)